arXiv:1609.06481v1 [math.NT] 21 Sep 2016

NEWFORMS OF HALF-INTEGRAL WEIGHT: THE MINUS SPACE COUNTERPART

EHUD MOSHE BARUCH AND SOMA PURKAIT

Abstract. We define a subspace of the space of holomorphic modular forms of weight $k+1/2$ and level $4M$ where M is odd and square-free. We show that this subspace is isomorphic under the Shimura-Niwa correspondence to the space of newforms of weight $2k$ and level $2M$ and that this is a Hecke isomorphism. The space we define is a proper subspace of the orthogonal complement of the Kohnen plus space if the Kohnen plus space is nonzero.

1. Introduction

Let M be odd and square-free and k be a positive integer. In a remarkable work, Niwa [7] comparing the traces of Hecke operators proved existence of Hecke isomorphism between the space of holomorphic cusp forms of weight $k+1/2$ on the congruence subgroup $\Gamma_0(4M)$ and the space of weight $2k$ cusp forms on $\Gamma_0(2M)$. In [4, 5] Kohnen defines plus space $S^+_{k+1/2}(4M)$, a subspace of the space $S_{k+1/2}(\Gamma_0(4M))$, which is given by a certain Fourier coefficient condition. Kohnen considers a new space $S^{+,\text{new}}_{k+1/2}(4M)$ inside his plus space and proves that this new subspace is Hecke isomorphic to $S^{\text{new}}_{2k}(\Gamma_0(M))$, the space of newforms of weight $2k$ and level M, giving a newform theory of half-integral weight.

In the case $M=1$ the Kohnen plus space is given as an eigenspace of a certain Hecke operator considered by Niwa [7]. Niwa's operator has two eigenvalues, one positive and one negative and the Kohnen plus space is the eigenspace of the positive eigenvalue. In this case the Kohnen plus space itself is the newspace $S^{+,\text{new}}_{k+1/2}(4)$ and is isomorphic as a Hecke module to $S_{2k}(\Gamma_0(1))$. In the case $M>1$, the Kohnen plus space can be again given as an eigenspace of a certain operator [5] and we shall see that this operator at level $4M$ is indeed an analogue of Niwa's operator at level 4.

The Kohnen isomorphism uses the above mentioned Niwa isomorphism between the full spaces of weight $k+1/2$, level $4M$ and weight $2k$, level $2M$. From Kohnen's results, it is clear that the Niwa map sends the Kohnen plus space to a subspace of old forms inside $S_{2k}(\Gamma_0(2M))$. Kohnen's map can be

Date: August 2016.

2010 *Mathematics Subject Classification.* Primary: 11F37; Secondary: 11F12, 11F70.

Key words and phrases. Hecke algebras, Half-integral weight forms, Niwa isomorphism, Kohnen plus space.

viewed as a composition of the Niwa map with an another map which sends an old form of level $2M$ from this image to its newform originator at level M. Our objective in this paper is to identify the subspace of half-integer weight forms which is sent by the Niwa map isomorphically onto the space of newforms for $\Gamma_0(2M)$. It is clear that this space will complement the Kohnen plus space. However, their sum will not give the whole space if the Kohnen plus space is nonzero.

In fact, our results are motivated by the results of Loke and Savin [6] who interpreted the Kohnen plus space in representation theory language. For the case $M = 1$, Loke and Savin defined another space of half-integer weight forms which they showed is "conjugate" to the Kohnen plus space. This means that it is an image of the Kohnen plus space by an invertible Hecke operator and is isomorphic to the Kohnen plus space as a Hecke module. We show that the Kohnen plus space and the space considered by Loke and Savin do not intersect and that their sum maps isomorphically to the space of old forms $S_{2k}^{\text{old}}(\Gamma_0(2))$ under the Niwa map. Hence it is natural to consider the orthogonal complement of the direct sum under the Petersson inner product, which we call as the minus space. We show that the minus space is mapped isomorphically under the Niwa map to $S_{2k}^{\text{new}}(\Gamma_0(2))$, the space of newforms on $\Gamma_0(2)$. We also characterize this space as a common eigenspace of two Hecke operators: the Niwa operator used by Kohnen to define the Kohnen plus space and a conjugate of the Niwa operator which was considered by Loke and Savin. The minus space is the intersection of the negative eigenspace of both operators. We normalize the negative eigenvalue to be -1 as in [2]. Our description of the minus space at level 4 is completely analogous to our description of the new space $S_{2k}^{\text{new}}(\Gamma_0(2))$ in [2] where we showed that $S_{2k}^{\text{new}}(\Gamma_0(2))$ is the common -1 eigenspace of two Hecke operators. To summarize the case of $M = 1$: We show that the space $S_{k+1/2}(\Gamma_0(4))$ decomposes into a direct sum of three spaces: The Kohnen plus space, a "conjugate" of the Kohnen plus space given by Loke and Savin and the minus space. The Kohnen plus space and its conjugate are indistinguishable as Hecke modules which is the same as saying that they are mapped under the Niwa map to "conjugate" spaces of old forms. The minus space is different as a Hecke module from both spaces.

In order to generalize this result for M odd and square-free we consider a certain p-adic Hecke algebra for every prime p dividing M. Our work follows that of Loke and Savin who studies a certain 2-adic Hecke algebra which allowed them to give a representation theoretic interpretation of the Kohnen plus space and to introduce the operator which is conjugate of Niwa's operator and the space which is "conjugate" to Kohnen's plus space. In our p-adic Hecke algebra, we consider two p-adic operators that give rise to conjugate classical Hecke operators which when we use along with Niwa's operator and it's conjugate allow us to define our minus space at level $4M$. We give two descriptions of the minus space: One description as an orthogonal complement of a certain sum of subspaces and another description as a common

-1 eigenspace of the Niwa operator, its conjugate and a pair of conjugate operators for each prime dividing M. This again is completely analogous to our description of the space of newforms of weight $2k$ for $\Gamma_0(2M)$ given in [2]. Our main result is that the minus space of weight $k+1/2$ at level $4M$ is isomorphic as a Hecke module to the space of newforms of weight $2k$ at level $2M$.

Our paper is divided as follows. We start with recalling Gelbart's theory of double cover of $\mathrm{SL}_2(\mathbb{Q}_p)$. For an odd prime p we set up definition of genuine Iwahori Hecke algebra of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ modulo $\overline{K_0^p(p)}$ and a genuine central character. In section 3 we describe this Hecke algebra with generators and relations. In section 4 we translate certain elements in our p-adic Hecke algebra to classical Hecke operators on $S_{k+1/2}(\Gamma_0(4M))$. In particular we obtain two classical operators, $\widetilde{Q}_p$ with eigenvalues p and -1 and an involution $\widetilde{W}_{p^2}$. We further consider $\widetilde{Q}'_p$ which is conjugate of $\widetilde{Q}_p$ by $\widetilde{W}_{p^2}$. We check that these operators are self-adjoint with respect to Petersson inner product. In section 5 we revisit Loke and Savin and study Kohnen's classical operator Q on $S_{k+1/2}(\Gamma_0(4M))$ which he uses to describe his plus space. We show that his operator Q comes from the 2-adic Hecke algebra considered by Loke and Savin, and is an analogue of Niwa's operator at level 4. Let $\widetilde{Q}'_2 := \left(\frac{2}{2k+1}\right) Q/\sqrt{2}$ and $\widetilde{Q}_2$ be conjugate of $\widetilde{Q}'_2$ by an involution $\widetilde{W}_4$. In section 6 we define our minus space $S^-_{k+1/2}(4M)$ and prove our main result:

Theorem. *Let $S^-_{k+1/2}(4M) \subseteq S_{k+1/2}(\Gamma_0(4M))$ be the common -1 eigenspace of operators $\widetilde{Q}_p$ and $\widetilde{Q}'_p$ for all primes p dividing $2M$. Then $S^-_{k+1/2}(4M)$ has a basis of eigenforms for all the operators T_{q^2} where q is a prime coprime to $2M$ and maps isomorphically under the Niwa map onto the space $S^{\mathrm{new}}_{2k}(\Gamma_0(2M))$.*

2. Review

Let p be any prime (including infinite place). The group $\mathrm{SL}_2(\mathbb{Q}_p)$ has a non-trivial central extension by $\mu_2 = \{\pm 1\}$:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mu_2 & \longrightarrow & \widetilde{\mathrm{SL}}_2(\mathbb{Q}_p) & \longrightarrow & \mathrm{SL}_2(\mathbb{Q}_p) \longrightarrow 1 \\ & & \{(I, \pm 1)\} & & (g, \pm 1) & \longmapsto & g \end{array}$$

We use the 2-cocycle described by Gelbart [3] to determine the double cover $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$. For $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Q}_p)$, define

$$\tau(g) = \begin{cases} c & \text{if } c \neq 0 \\ d & \text{if } c = 0 \end{cases};$$

if $p=\infty$, set $s_p(g)=1$ while for a finite prime p

$$s_p(g)=\begin{cases}(c,d)_p & \text{if } cd\neq 0 \text{ and } \mathrm{ord}_p(c) \text{ is odd}\\ 1 & \text{else.}\end{cases}$$

Define the 2-cocycle σ_p on $\mathrm{SL}_2(\mathbb{Q}_p)$ as follows:

$$\sigma_p(g,h)=(\tau(gh)\tau(g),\tau(gh)\tau(h))_p\, s_p(g)s_p(h)s_p(gh).$$

Then the double cover $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ is the set $\mathrm{SL}_2(\mathbb{Q}_p)\times\mu_2$ with the group law:

$$(g,\ \epsilon_1)(h,\ \epsilon_2)=(gh,\ \epsilon_1\epsilon_2\sigma_p(g,h)).$$

For any subgroup H of $\mathrm{SL}_2(\mathbb{Q}_p)$, we shall denote by $\overline{H}$ the complete inverse image of H in $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$.

We consider the following subgroups of $\mathrm{SL}_2(\mathbb{Z}_p)$:

$$K_0^p(p^n)=\left\{\begin{pmatrix}a & b\\ c & d\end{pmatrix}\in \mathrm{SL}_2(\mathbb{Z}_p)\ :\ c\in p^n\mathbb{Z}_p\right\},$$

$$K_1^p(p^n)=\left\{\begin{pmatrix}a & b\\ c & d\end{pmatrix}\in \mathrm{SL}_2(\mathbb{Z}_p)\ :\ c\in p^n\mathbb{Z}_p,\ a\equiv 1 \pmod{p^n\mathbb{Z}_p}\right\}.$$

By [3, Proposition 2.8] for odd primes p, $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ splits over $\mathrm{SL}_2(\mathbb{Z}_p)$. Thus $\overline{\mathrm{SL}_2(\mathbb{Z}_p)}$ is isomorphic to direct product of $\mathrm{SL}_2(\mathbb{Z}_p)$ and μ_2 and, $\overline{K_0^p(p)}$ is direct product of $K_0^p(p)$ and μ_2. Hence it follows from [3, Corollary 2.13] that the center M_p of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ is direct product $\{\pm I\}\times\mu_2$. Thus any genuine central character is given by a a non-trivial character of $\mu_2\times\mu_2$.

However $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_2)$ does not split over $\mathrm{SL}_2(\mathbb{Z}_2)$ but instead splits over subgroup

$$K_1^2(4)=\left\{\begin{pmatrix}a & b\\ c & d\end{pmatrix}\in \mathrm{SL}_2(\mathbb{Z}_2)\ :\ c\equiv 0,\ a\equiv 1 \pmod{4\mathbb{Z}_2}\right\}.$$

In this case the center M_2 of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_2)$ is a cyclic group of order 4 generated by $(-I,1)$ and so a genuine central character is given by a sending $(-I,1)$ to a primitive fourth root of unity. One can check that $\overline{K_0^2(4)}$ is a direct product $K_1^2(4)\times M$ (see [6]).

Loke and Savin [6] considered a genuine Hecke algebra for $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_2)$ corresponding to $\overline{K_0^2(4)}$ and a genuine central character. We shall study genuine Hecke algebras for $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ corresponding to $\overline{K_0^p(p)}$ and a given genuine character of M_p for general odd prime p.

Before moving to the next section we set up notation: For $s\in\mathbb{Q}_p$, $t\in\mathbb{Q}_p^\times$ we shall consider the following elements of $\mathrm{SL}_2(\mathbb{Q}_p)$:

$$x(s)=\begin{pmatrix}1 & s\\ 0 & 1\end{pmatrix},\ y(s)=\begin{pmatrix}1 & 0\\ s & 1\end{pmatrix},\ w(t)=\begin{pmatrix}0 & t\\ -t^{-1} & 0\end{pmatrix},\ h(t)=\begin{pmatrix}t & 0\\ 0 & t^{-1}\end{pmatrix}.$$

Let $N = \{(x(s), \epsilon) : s \in \mathbb{Q}_p,\ \epsilon = \pm 1\}$, $T = \{(h(t), \epsilon) : t \in \mathbb{Q}_p^\times,\ \epsilon = \pm 1\}$ and $\bar{N} = \{(y(s), \epsilon) : s \in \mathbb{Q}_p,\ \epsilon = \pm 1\}$ be the subgroups of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$. For any subgroup S of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$, let $N^S = N \cap S$, $T^S = T \cap S$ and $\bar{N}^S = \bar{N} \cap S$.

In the next section we will need some lemmas on cocycle multiplication, we note them in the appendix.

3. Iwahori Hecke Algebra of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ modulo $\overline{K_0^p(p)}$, p odd.

Fix an odd prime p. In this section we shall denote $K_0^p(p)$ by simply K_0. Let γ be a character of K_0 such that it is trivial on $K_1^p(p)$. Since $\frac{K_0^p(p)}{K_1^p(p)} \cong (\mathbb{Z}_p/p\mathbb{Z}_p)^\times$ by $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto d \pmod{p\mathbb{Z}_p}$, we can define γ by a character of $(\mathbb{Z}/p\mathbb{Z})^\times$. We shall use the symbol γ to also denote a genuine character on $\overline{K}_0$ by defining $\gamma(A, \epsilon) = \epsilon\gamma(A)$ for $A \in K_0$.

We shall consider "genuine " Iwahori Hecke algebra of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ with respect to $\overline{K}_0$ and central character γ.

Let $C_c^\infty(\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p))$ denote the space of locally constant, compactly supported complex-valued functions on G. Let $H(\overline{K}_0, \gamma)$ be the subalgebra defined as follows:

$$\{f \in C_c^\infty(\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)) : f(\tilde{k}\tilde{g}\tilde{k}') = \overline{\gamma}(\tilde{k})\overline{\gamma}(\tilde{k}')f(\tilde{g}) \text{ for } \tilde{g} \in \widetilde{\mathrm{SL}}_2(\mathbb{Q}_p),\ \tilde{k},\ \tilde{k}' \in \overline{K}_0\}.$$

Then $H(\overline{K}_0, \gamma)$ is a $\mathbb{C}$-algebra under convolution which, for any $f_1, f_2 \in H(\overline{K}_0, \gamma)$, is defined by

$$f_1 * f_2(\tilde{h}) = \int_{\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)} f_1(\tilde{g}) f_2(\tilde{g}^{-1}\tilde{h}) d\tilde{g} = \int_{\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)} f_1(\tilde{h}\tilde{g}) f_2(\tilde{g}^{-1}) d\tilde{g},$$

where $d\tilde{g}$ is the Haar measure on $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ such that the measure of $\overline{K}_0$ is one.

We want to describe $H(\overline{K}_0, \gamma)$ using generators and relations.

Lemma 3.1. *A complete set of representatives for the double cosets of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ mod $\overline{K}_0$ are given by $(h(p^n), 1)$, $(w(p^{-n}), 1)$ where n varies over integers.*

We say that $H(\overline{K}_0, \gamma)$ is supported on $\tilde{g} \in \widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ if there exists $f \in H(\overline{K}_0, \gamma)$ such that $f(\tilde{g}) \neq 0$.

Lemma 3.2. *Let $K_{\tilde{g}} = \overline{K}_0 \cap \tilde{g}\overline{K}_0\tilde{g}^{-1}$. Then $H(\overline{K}_0, \gamma)$ is supported on $\tilde{g}$ if and only if for every $\tilde{k} \in K_{\tilde{g}}$ we have $\gamma([\tilde{k}^{-1},\ \tilde{g}^{-1}]) = 1$.*

Lemma 3.3. *The function $\alpha_{\tilde{g}} : K_{\tilde{g}} \longrightarrow \mathbb{C}$ defined by $\alpha_{\tilde{g}}(\tilde{k}) = \gamma([\tilde{k}^{-1},\ \tilde{g}^{-1}])$ is a character of $K_{\tilde{g}}$.*

Remark 1. *In the above lemmas one can replace $\overline{K}_0$ by any subgroup S of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ and γ by any character of such S. Further p can be any prime.*

3.1. **Support of $H(\overline{K}_0,\gamma)$.** Let p be an odd prime. Fix an integer n. Let $A := h(p^n)$ and $\tilde{A} := (A,\epsilon_1)$. We shall show that $H(\overline{K}_0,\gamma)$ is supported on $\tilde{A}$. We have

$$K_{\tilde{A}} := \Big\{ \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \pm 1 \right) \in \overline{SL}_2(\mathbb{Z}_p) \ : \mathrm{ord}_p(c) \geq \max\{-2n+1,1\}, \\ \mathrm{ord}_p(b) \geq \max\{2n,0\} \Big\}.$$

As in Lemma A.4 we check that $K_{\tilde{A}}$ has a triangular decomposition $K_{\tilde{A}} = N^{K_{\tilde{A}}} T^{K_{\tilde{A}}} \bar{N}^{K_{\tilde{A}}}$ where $T^{K_{\tilde{A}}} = T^{\overline{K}_0}$,

$$N^{K_{\tilde{A}}} := \{(x(s),\pm 1) : \mathrm{ord}_p(s) \geq \max\{2n,0\}\},$$

$$\bar{N}^{K_{\tilde{A}}} := \{(y(t),\pm 1) : \mathrm{ord}_p(t) \geq \max\{-2n+1,1\}\},$$

By Lemma 3.2 and 3.3, it is enough to check that the value of γ on the commutator $[(B,\ \epsilon_2)^{-1},\ (A,\ \epsilon_1)^{-1}]$ is 1 for any $(B,\ \epsilon_2)$ in $N^{K_{\tilde{A}}}$, $T^{K_{\tilde{A}}}$ and $\bar{N}^{K_{\tilde{A}}}$ respectively.

By Lemma A.3, for $B = (x(s),\epsilon_2) \in N^{K_{\tilde{A}}}$, we get $[(B,\ \epsilon_2)^{-1},\ (A,\ \epsilon_1)^{-1}] = (\begin{pmatrix} 1 & sp^{-2n} - s \\ 0 & 1 \end{pmatrix}, 1)$; for $B = (h(u),\epsilon_2) \in T^{K_{\tilde{A}}}$, $[(B,\ \epsilon_2)^{-1},\ (A,\ \epsilon_1)^{-1}] = (I,1)$; and for $B = (y(t),\epsilon_2) \in N^{K_{\tilde{A}}}$, we get that $[(B,\ \epsilon_2)^{-1},\ (A,\ \epsilon_1)^{-1}] = (\begin{pmatrix} 1 & 0 \\ (p^{2n}-1)t & 1 \end{pmatrix}, 1)$. Since each of them belong to $K_1^p(p) \times \{1\}$, we are done.

Next let $A := w(p^{-n})$. We show that $H(\overline{K}_0,\gamma)$ is supported on $\tilde{A} = (A,\epsilon_1)$ provided either $\gamma(u^2) = 1$ for all units u in $\mathbb{Z}_p$. In this case we have

$$K_{\tilde{A}} := \Big\{ \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \pm 1 \right) \in \overline{SL}_2(\mathbb{Z}_p) \ : \mathrm{ord}_p(c) \geq \max\{2n,1\}, \\ \mathrm{ord}_p(b) \geq \max\{-2n+1,0\} \Big\}.$$

We again check that $K_{\tilde{A}}$ has a triangular decomposition $K_{\tilde{A}} = N^{K_{\tilde{A}}} T^{K_{\tilde{A}}} \bar{N}^{K_{\tilde{A}}}$ and here $T^{K_{\tilde{A}}} = T^{\overline{K}_0}$,

$$N^{K_{\tilde{A}}} := \{(x(s),\pm 1) : \mathrm{ord}_p(s) \geq \max\{-2n+1,0\}\},$$

$$\bar{N}^{K_{\tilde{A}}} := \{(y(t),\pm 1) : \mathrm{ord}_p(t) \geq \max\{2n,1\}\}.$$

By Lemma A.3, for $B = (x(s),\epsilon_2) \in N^{K_{\tilde{A}}}$, we get $[(B,\ \epsilon_2)^{-1},\ (A,\ \epsilon_1)^{-1}] = (\begin{pmatrix} 1+s^2p^{2n} & -s \\ -sp^{2n} & 1 \end{pmatrix}, 1)$, so γ takes value 1 on this commutator. In the case $B = (y(t),\epsilon_2) \in N^{K_{\tilde{A}}}$, we have

$$B^{-1}A^{-1}BA = \begin{pmatrix} 1 & -p^{-2n}t \\ -t & 1+p^{-2n}t^2 \end{pmatrix}, \text{ where } \mathrm{ord}_p(t) \geq \max\{2n,1\},$$

so $s_p(B^{-1}A^{-1}BA) = 1$ if either $-t(1+p^{-2n}t^2) = 0$ or $\mathrm{ord}_p(t)$ is even. Assume that $-t(1+p^{-2n}t^2) \neq 0$ and $\mathrm{ord}_p(t)$ is odd. Then $s_p(B^{-1}A^{-1}BA) =$

$\left(-t, 1+p^{-2n}t^2\right)_p = \left(-p, 1+p^{-2n}t^2\right)_p$. Let $u = 1+p^{-2n}t^2$. Since $\mathrm{ord}_p(t) \geq \max\{2n, 1\}$, we have $u \equiv 1 \pmod{p\mathbb{Z}_p}$. Hence $s_p(B^{-1}A^{-1}BA) = (-p, u)_p = \left(\frac{u}{p}\right) = 1$. So in this case also γ takes value 1.

For $B = (h(u), \epsilon_2) \in T^{K_{\tilde{A}}}$, $[(B,\ \epsilon_2)^{-1},\ (A,\ \epsilon_1)^{-1}] = (\begin{pmatrix} 1/u^2 & 0 \\ 0 & u^2 \end{pmatrix}, 1)$, so $\gamma([(B,\ \epsilon_2)^{-1},\ (A,\ \epsilon_1)^{-1}]) = \gamma(u^2)$.

Thus if $\gamma(u^2) = 1$ for all units u in $\mathbb{Z}_p$ then $H(\overline{K}_0, \gamma)$ is supported on $(w(p^{-n}), \epsilon)$. In particular this holds if our choice of γ is quadratic.

Proposition 3.4. *If γ is a quadratic character then $H(\overline{K}_0, \gamma)$ is supported on the double cosets of $\overline{K}_0$ represented by $(h(p^n), 1)$ and $(w(p^{-n}), 1)$ where n varies over integers.*

We have the following decomposition that can be obtained using triangular decomposition of K_0.

Lemma 3.5. *1. For $n \geq 0$,*

$$K_0 h(p^n) K_0 = \bigcup_{s \in \mathbb{Z}_p/p^{2n}\mathbb{Z}_p} x(s)h(p^n)K_0 = \bigcup_{s \in \mathbb{Z}_p/p^{2n}\mathbb{Z}_p} K_0 h(p^n) y(ps).$$

2. For $n \geq 1$,

$$K_0 h(p^{-n}) K_0 = \bigcup_{s \in \mathbb{Z}_p/p^{2n}\mathbb{Z}_p} y(ps)h(p^{-n})K_0 = \bigcup_{s \in \mathbb{Z}_p/p^{2n}\mathbb{Z}_p} K_0 h(p^{-n}) x(s).$$

3. For $n \geq 1$,

$$K_0 w(p^{-n}) K_0 = \bigcup_{s \in \mathbb{Z}_p/p^{2n-1}\mathbb{Z}_p} y(ps)w(p^{-n})K_0 = \bigcup_{s \in \mathbb{Z}_p/p^{2n-1}\mathbb{Z}_p} K_0 w(p^{-n}) y(ps).$$

4. For $n \geq 0$,

$$K_0 w(p^{n}) K_0 = \bigcup_{s \in \mathbb{Z}_p/p^{2n+1}\mathbb{Z}_p} x(s)w(p^{n})K_0 = \bigcup_{s \in \mathbb{Z}_p/p^{2n+1}\mathbb{Z}_p} K_0 w(p^{n}) x(s).$$

We note the following well-known lemmas that will be useful in computing convolutions.

Lemma 3.6. *Let $f_1,\ f_2 \in H(\overline{K}_0, \gamma)$ such that f_1 is supported on $\overline{K}_0 \tilde{x} \overline{K}_0 = \bigcup_{i=1}^m \tilde{\alpha}_i \overline{K}_0$ and f_2 is supported on $\overline{K}_0 \tilde{y} \overline{K}_0 = \bigcup_{j=1}^n \tilde{\beta}_i \overline{K}_0$. Then*

$$f_1 * f_2(\tilde{h}) = \sum_{i=1}^{m} f_1(\tilde{\alpha}_i) f_2(\tilde{\alpha}_i^{-1}\tilde{h})$$

where the nonzero summands are precisely for those i for which there exist a j such that $\tilde{h} \in \tilde{\alpha}_i \tilde{\beta}_j \overline{K}_0$.

For $\tilde{g} \in \widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ let $\mu(\tilde{g})$ denotes the number of disjoint left (right) $\overline{K}_0$ cosets in the decomposition of double coset $\overline{K}_0 \tilde{g} \overline{K}_0$.

Lemma 3.7. *Let $\tilde{g},\ \tilde{h} \in \widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ be such that $\mu(\tilde{g})\mu(\tilde{h}) = \mu(\tilde{g}\tilde{h})$. Let f_1 and $f_2 \in H(\overline{K}_0, \gamma)$ are respectively supported on $\overline{K}_0\tilde{g}\overline{K}_0$ and $\overline{K}_0\tilde{h}\overline{K}_0$. Then $f_1 * f_2$ is precisely supported on $\overline{K}_0\tilde{g}\tilde{h}\overline{K}_0$ and $f_1 * f_2(\tilde{g}\tilde{h}) = f_1(\tilde{g})f_2(\tilde{h})$.*

3.2. **Generators and relations.** We continue with the notation as in the previous subsection. We shall extend the character γ of $\overline{K}_0$ to the subgroup T of $\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$ and then to the normalizer group $N_{\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)}(T)$ which is generated by elements $(h(t), \epsilon)$, $(w(t), \epsilon)$ for $t \in \mathbb{Q}_p^\times$. We note the following relations:

$$(h(s),1)(h(t),1) = (h(st), (s,t)_p), \quad (w(s),1)(w(t),1) = (h(-st^{-1}), (s,t)_p),$$

$$(h(s),1)(w(t),1) = (w(st), (s,-t)_p), \ \ (w(s),1)(h(t),1) = (w(st^{-1}), (-s,t)_p).$$

Let $\varepsilon_p = 1$ or $(-1)^{1/2}$ depending on whether $p \equiv 1$ or $3 \pmod 4$. Thus $\varepsilon_p^2 = \left(\frac{-1}{p}\right)$. Define

$$\gamma((h(p^n),1)) = \begin{cases} 1 & \text{if } n \text{ is even} \\ \varepsilon_p & \text{if } n \text{ is odd.} \end{cases}$$

For a general $t = p^n u \in \mathbb{Q}_p$ where $n \in \mathbb{Z}$ and u is a unit in $\mathbb{Z}_p$, we use the relation $(h(t),1) = (h(p^n),1)(h(u),1)(I, (p^n,u)_p)$ to define

$$\gamma((h(t),1)) = \begin{cases} 1 & \text{if } n \text{ is even} \\ \varepsilon_p \left(\frac{u}{p}\right) & \text{if } n \text{ is odd.} \end{cases}$$

One can check that γ indeed extends to a character on T.

We now extend the character to the normalizer $N_{\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)}(T)$ by defining $\gamma((w(1),1)) = 1$ and extend it using the following relation

$$(w(t),1) = (h(t),1)(w(1),1)(I, \left(-1, t^{-1}\right)_p).$$

For $t = p^n u$ as above we have $\left(-1, t^{-1}\right)_p = 1$ for n even and $\left(-1, t^{-1}\right)_p = (-1, pu)_p = \left(\frac{-1}{p}\right)$ if n is odd, hence

$$\gamma((w(t),1)) = \begin{cases} 1 & \text{if } n \text{ is even} \\ \varepsilon_p \left(\frac{-u}{p}\right) & \text{if } n \text{ is odd.} \end{cases}$$

We define the following elements of $H(\overline{K}_0, \gamma)$ supported respectively on the double cosets of $(h(p^n),1)$ and $(w(p^{-n}),1)$: For $n \in \mathbb{Z}$, and $\tilde{k}, \tilde{k}' \in \overline{K}_0$,

$$X_{(h(p^n),1)}(\tilde{k}(h(p^n),1)\tilde{k}') = \overline{\gamma}(\tilde{k})\overline{\gamma}((h(p^n),1))\overline{\gamma}(\tilde{k}'),$$

$$X_{(w(p^{-n}),1)}(\tilde{k}(w(p^{-n}),1)\tilde{k}') = \overline{\gamma}(\tilde{k})\overline{\gamma}((w(p^{-n}),1))\overline{\gamma}(\tilde{k}').$$

Let $\mathcal{T}_n = X_{(h(p^n),1)}$ and $\mathcal{U}_n = X_{(w(p^{-n}),1)}$. Then Proposition 3.4 implies that $\mathcal{T}_n$ and $\mathcal{U}_n$ forms a $\mathbb{C}$-basis for $H(\overline{K}_0, \gamma)$.

Proposition 3.8. *We have following relations:*

(1) If $mn \geq 0$ then $\mathcal{T}_m * \mathcal{T}_n = \mathcal{T}_{m+n}$.

(2) For $n \geq 0$, $\mathcal{U}_1 * \mathcal{T}_n = \mathcal{U}_{n+1}$.
(3) For $n \geq 0$, $\mathcal{T}_{-n} * \mathcal{U}_1 = \mathcal{U}_{n+1}$.
(4) For $n \geq 0$, $\mathcal{U}_0 * \mathcal{T}_{-n} = \mathcal{U}_{-n}$.
(5) For $n \geq 0$, $\mathcal{T}_n * \mathcal{U}_0 = \mathcal{U}_{-n}$.
(6) For $n \geq 1$, $\mathcal{U}_0 * \mathcal{U}_n = \overline{\gamma}(-1) \cdot \mathcal{T}_n$.
(7) For $n \geq 1$, $\mathcal{U}_n * \mathcal{U}_0 = \overline{\gamma}(-1) \cdot \mathcal{T}_{-n}$.

Proof. We prove (1) and (7). The rest are similar. For (1) let $mn \geq 0$. We may assume both $m,\ n \geq 0$. It follows from Lemma 3.5 and 3.7 that $\mathcal{T}_m * \mathcal{T}_n$ is precisely supported on the double coset $\overline{K}_0(h(p^{n+m}),1)\overline{K}_0$ and that

$$\mathcal{T}_m * \mathcal{T}_n((h(p^m),1)(h(p^n),1)) = \mathcal{T}_m((h(p^m),1))\mathcal{T}_n((h(p^n),1)).$$

Let m and n both be even. Then $(h(p^m),1)(h(p^n),1) = (h(p^{n+m}),1)$ and so

$$\mathcal{T}_m * \mathcal{T}_n((h(p^{n+m}),1)) = \mathcal{T}_m((h(p^m),1))\mathcal{T}_n((h(p^n),1)) =$$
$$= \overline{\gamma}((h(p^m),1))\overline{\gamma}((h(p^n),1)) = 1 = \mathcal{T}_{m+n}((h(p^{n+m}),1)),$$

hence $\mathcal{T}_m * \mathcal{T}_n = \mathcal{T}_{m+n}$. Next suppose both m and n are odd, so $m+n$ is even. Then $(h(p^m),1)(h(p^n),1) = (h(p^{n+m}),1)(I,(p,p)_p)$ and so

$$\mathcal{T}_m * \mathcal{T}_n((h(p^{n+m}),1)) = \overline{\gamma}((I,(p,p)_p))\mathcal{T}_m((h(p^m),1))\mathcal{T}_n((h(p^n),1)) =$$
$$= \left(\frac{-1}{p}\right)\overline{\gamma}((h(p^m),1))\overline{\gamma}((h(p^n),1)) = \left(\frac{-1}{p}\right)\overline{\varepsilon_p^2} = 1 = \mathcal{T}_{m+n}((h(p^{n+m}),1)).$$

Now suppose m is odd and n is even (or vice versa), so $m+n$ is odd. In this case $(h(p^m),1)(h(p^n),1) = (h(p^{n+m}),1)$ and so

$$\mathcal{T}_m * \mathcal{T}_n((h(p^{n+m}),1)) = \overline{\varepsilon_p} = \mathcal{T}_{m+n}((h(p^{n+m}),1))$$

and we are done.

For (7), let $n \geq 1$. As before using Lemma 3.5 and 3.7 we know that $\mathcal{U}_n * \mathcal{U}_0$ is supported on the double coset $\overline{K}_0(h(p^{-n}),1)\overline{K}_0$ and that

$$\mathcal{U}_n * \mathcal{U}_0((w(p^{-n}),1)(w(1),1)) = \mathcal{U}_n((w(p^{-n}),1))\mathcal{U}_0((w(1),1)).$$

We have $(w(p^{-n}),1)(w(1),1) = (h(p^{-n}),1)(-I,(p^{-n},-1)_p)$ and so

$$\overline{\gamma}(-1)\ \mathcal{U}_n * \mathcal{U}_0((h(p^{-n}),1)) = \left(p^{-n},-1\right)_p \mathcal{U}_n((w(p^{-n}),1))\mathcal{U}_0((w(1),1)) =$$
$$= \begin{cases} \left(\frac{-1}{p}\right)\overline{\varepsilon_p}\left(\frac{-1}{p}\right) = \overline{\varepsilon_p} & \text{if } n \text{ is odd} \\ 1 & \text{if } n \text{ is even} \end{cases} = \mathcal{T}_{-n}((h(p^{-n}),1)),$$

and thus $\mathcal{U}_n * \mathcal{U}_0 = \overline{\gamma}(-1) \cdot \mathcal{T}_{-n}$. □

We will consider two choices for γ as a character of $(\mathbb{Z}/p\mathbb{Z})^*$, either γ is trivial or γ is given by the Kronecker symbol $\gamma = \left(\frac{\cdot}{p}\right)$. Then we have following proposition.

Proposition 3.9. *We have following relations.*

(1) $\mathcal{U}_0^2 = \begin{cases} (p-1)\mathcal{U}_0 + p & \text{if } \gamma \text{ is trivial} \\ \left(\frac{-1}{p}\right)p & \text{if } \gamma = \left(\frac{\cdot}{p}\right). \end{cases}$

(2) $\mathcal{U}_1^2 = \begin{cases} p & \text{if } \gamma \text{ is trivial} \\ \varepsilon_p(p-1)U_1 + \left(\frac{-1}{p}\right)p & \text{if } \gamma = \left(\frac{\cdot}{p}\right). \end{cases}$

(3) If γ *is trivial, then* $\mathcal{T}_1 * \mathcal{U}_1 = p\,\mathcal{U}_0$ *and* $\mathcal{T}_{-1} = 1/p \cdot \mathcal{U}_1 * \mathcal{T}_1 * \mathcal{U}_1$.

Proof. For (1) we use Lemma 3.6 to check that $\mathcal{U}_0 * \mathcal{U}_0$ is at most supported on the double cosets $\overline{K}_0$ and $\overline{K}_0(w(1),1)\overline{K}_0$. Thus we need to only compute the values of $\mathcal{U}_0^2$ at $(I,1)$ and $(w(1),1)$. Using Lemma 3.5 and 3.6 we have

$$\begin{aligned}
\mathcal{U}_0 * \mathcal{U}_0((I,1)) &= \sum_{s=0}^{p-1} \mathcal{U}_0((x(s),1)(w(1),1))\mathcal{U}_0((w(1),1)^{-1}(x(s),1)^{-1}) \\
&= \sum_{s=0}^{p-1} \mathcal{U}_0((w(1),1))\mathcal{U}_0((w(-1),1)(x(-s),1)) \\
&= \sum_{s=0}^{p-1} \mathcal{U}_0((h(-1),1)(w(1),1)(x(-s),1)) \\
&= \sum_{s=0}^{p-1} \overline{\gamma}(-1) = \begin{cases} p & \text{if } \gamma \text{ is trivial} \\ \left(\frac{-1}{p}\right)p & \text{if } \gamma = \left(\frac{\cdot}{p}\right). \end{cases}
\end{aligned}$$

Similarly, we get that $\mathcal{U}_0 * \mathcal{U}_0((w(1),1)) =$

$$\begin{aligned}
&= \sum_{s=0}^{p-1} \mathcal{U}_0((x(s),1)(w(1),1))\mathcal{U}_0((w(1),1)^{-1}(x(s),1)^{-1}(w(1),1)) \\
&= \sum_{s=0}^{p-1} \mathcal{U}_0((\begin{pmatrix} 0 & -1 \\ 1 & -s \end{pmatrix},1)(w(1),1))) = \sum_{s=0}^{p-1} \mathcal{U}_0((y(s),1)) \\
&= \sum_{s=1}^{p-1} \mathcal{U}_0((y(s),1)) \quad \text{since } \mathcal{U}_0((I,1)) = 0.
\end{aligned}$$

It is easy to check that for $1 \le s \le p-1$

$$(y(s),1) = (\begin{pmatrix} 1 & 1/s \\ 0 & 1 \end{pmatrix},1)(w(1),1)(\begin{pmatrix} -s & -1 \\ 0 & -1/s \end{pmatrix},1) \in \overline{K}_0(w(1),1)\overline{K}_0$$

and hence $\mathcal{U}_0 * \mathcal{U}_0((w(1),1)) =$

$$= \sum_{s=1}^{p-1} \overline{\gamma}(-1/s) = \sum_{s=1}^{p-1} \gamma(s) = \begin{cases} p-1 & \text{if } \gamma \text{ is trivial} \\ \sum_{s=1}^{p-1}\left(\frac{s}{p}\right) = 0 & \text{if } \gamma = \left(\frac{\cdot}{p}\right). \end{cases}$$

Thus if we write $\mathcal{U}_0^2 = c_1\mathcal{U}_0 + c_2$, we get that

$$c_1 = \begin{cases} p-1 & \text{if } \gamma \text{ trivial} \\ 0 & \text{if } \gamma = \left(\frac{\cdot}{p}\right), \end{cases} \quad c_2 = \begin{cases} p & \text{if } \gamma \text{ trivial} \\ \left(\frac{-1}{p}\right)p & \text{if } \gamma = \left(\frac{\cdot}{p}\right). \end{cases}$$

Now we prove (2). Again using Lemma 3.6 we see that $\mathcal{U}_1 * \mathcal{U}_1$ is at most supported on the double cosets $\overline{K}_0$ and $\overline{K}_0(w(p^{-1}),1)\overline{K}_0$. So we need to

find the values of $\mathcal{U}_1^2$ at $(I,1)$ and $(w(p^{-1}),1)$. Using Lemma 3.5 and 3.6,

$$\begin{aligned}
\mathcal{U}_1 * \mathcal{U}_1((I,1)) &= \sum_{s=0}^{p-1} \mathcal{U}_1((y(ps),1)(w(p^{-1}),1))\mathcal{U}_1((w(p^{-1}),1)^{-1}(y(ps),1)^{-1}) \\
&= \sum_{s=0}^{p-1} \mathcal{U}_1((w(p^{-1}),1))\mathcal{U}_1((w(-p^{-1}),1)(y(-ps),1)) \\
&= \sum_{s=0}^{p-1} \overline{\varepsilon_p}\left(\frac{-1}{p}\right) \mathcal{U}_1((h(-1),(-p,-1)_p)(w(p^{-1}),1)) \\
&= \sum_{s=0}^{p-1} \overline{\varepsilon_p}\left(\frac{-1}{p}\right) \gamma(-1)\left(\frac{-1}{p}\right) \overline{\varepsilon_p}\left(\frac{-1}{p}\right) \\
&= \gamma(-1)p = \begin{cases} p & \text{if } \gamma \text{ trivial} \\ \left(\frac{-1}{p}\right) p & \text{if } \gamma = \left(\frac{\cdot}{p}\right). \end{cases}
\end{aligned}$$

Finally, we have $\mathcal{U}_1 * \mathcal{U}_1((w(p^{-1}),1)) =$

$$\begin{aligned}
&= \sum_{s=0}^{p-1} \mathcal{U}_1((y(ps),1)(w(p^{-1}),1))\mathcal{U}_1((w(-p^{-1}),1)(y(-ps),1)(w(p^{-1}),1)) \\
&= \sum_{s=0}^{p-1} \overline{\varepsilon_p}\left(\frac{-1}{p}\right) \mathcal{U}_1((\begin{pmatrix} s & -p^{-1} \\ p & 0 \end{pmatrix}, \left(p^2, -p^2 s\right)_p)(w(p^{-1}),1))) \\
&= \sum_{s=0}^{p-1} \overline{\varepsilon_p}\left(\frac{-1}{p}\right) \mathcal{U}_1((x(s/p),(p,-p)_p)) = \sum_{s=1}^{p-1} \overline{\varepsilon_p}\left(\frac{-1}{p}\right) \mathcal{U}_1((x(s/p),1)).
\end{aligned}$$

Now we check that for $1 \le s \le p-1$

$$(x(s/p),1)(I,\begin{pmatrix} s/p \end{pmatrix}) = (\begin{pmatrix} s & 0 \\ p & 1/s \end{pmatrix},1)(w(p^{-1}),1)(\begin{pmatrix} 1 & 0 \\ p/s & 1 \end{pmatrix},1)$$

and so

$$\begin{aligned}
\mathcal{U}_1 * \mathcal{U}_1((w(p^{-1}),1)) &= \sum_{s=1}^{p-1} \overline{\varepsilon_p}\left(\frac{-1}{p}\right)\left(\frac{s}{p}\right) \overline{\gamma}(1/s) \overline{\varepsilon_p}\left(\frac{-1}{p}\right) \\
&= \sum_{s=1}^{p-1} \left(\frac{-s}{p}\right) \overline{\gamma}(1/s) \\
&= \begin{cases} \sum_{s=1}^{p-1} \left(\frac{-s}{p}\right) = 0 & \text{if } \gamma \text{ trivial} \\ \sum_{s=1}^{p-1} \left(\frac{-s}{p}\right)\left(\frac{s^{-1}}{p}\right) = \left(\frac{-1}{p}\right)(p-1) & \text{if } \gamma = \left(\frac{\cdot}{p}\right). \end{cases}
\end{aligned}$$

Thus if we write $\mathcal{U}_1^2 = c_1\mathcal{U}_1 + c_2$, we get that

$$c_1 = \begin{cases} 0 & \text{if } \gamma \text{ trivial} \\ \varepsilon_p(p-1) & \text{if } \gamma = \left(\frac{\cdot}{p}\right), \end{cases} \quad c_2 = \begin{cases} p & \text{if } \gamma \text{ trivial} \\ \left(\frac{-1}{p}\right) p & \text{if } \gamma = \left(\frac{\cdot}{p}\right). \end{cases}$$

For (3) let γ be a trivial character. From Prop 3.8(6), we have $\mathcal{U}_0 * \mathcal{U}_1 = \mathcal{T}_1$. Right multiplication by $\mathcal{U}_1$ on both sides and using (2) above gives $\mathcal{T}_1 * \mathcal{U}_1 = p\ \mathcal{U}_0$. Further using Prop 3.8(7) we get that $\mathcal{T}_{-1} = \mathcal{U}_1 * \mathcal{U}_0 = 1/p \cdot \mathcal{U}_1 * \mathcal{T}_1 * \mathcal{U}_1$. □

Remark 2. *We compare the p-adic operator $\mathcal{U}_1$ with Ueda's classical operator Y_p* [11, Proposition 1.27] *which satisfies a similar relation. In particular if we consider operator $\mathcal{U}_1' = \overline{\varepsilon_p}\mathcal{U}_1$, then in the case γ is trivial we have*

$$(\mathcal{U}_1')^2 = (\overline{\varepsilon_p}\mathcal{U}_1)^2 = \varepsilon_p^2 p = \left(\frac{-1}{p}\right) p,$$

while in the case $\gamma = \left(\frac{\cdot}{p}\right)$ we have

$$(\mathcal{U}_1')^2 = (\overline{\varepsilon_p}\mathcal{U}_1)^2 = \overline{\varepsilon_p}^2 \left(\varepsilon_p(p-1)\mathcal{U}_1 + p\left(\frac{-1}{p}\right)\right) = (p-1)\mathcal{U}_1' + p.$$

Thus $\mathcal{U}_1'$ satisfies exactly the same relations as the operator Y_p.

Theorem 1. *The "genuine" Iwahori Hecke algebra $H(\overline{K_0^p(p)}, \gamma)$ for γ trivial or $\left(\frac{\cdot}{p}\right)$ is generated as an $\mathbb{C}$-algebra by $\mathcal{U}_0$ and $\mathcal{U}_1$ with the defining relations given by above proposition.*

Proof. We let $\mathcal{A}$ be an abstract algebra generated by $\tilde{\mathcal{U}_0}$ and $\tilde{\mathcal{U}_1}$ with defining relations as (1) and (2) of Proposition 3.9. We have a homomorphism from $\mathcal{A}$ to $H(\gamma)$ mapping $\tilde{\mathcal{U}_0}$ to $\mathcal{U}_0$ and $\tilde{\mathcal{U}_1}$ to $\mathcal{U}_1$. It follows from Proposition 3.8 that this homomorphism is onto. We let M be the kernel of this homomorphism. Using relations (1) and (2) it follows that M is a linear combination of words of the form $\tilde{\mathcal{U}_0}\tilde{\mathcal{U}_1}\tilde{\mathcal{U}_0}$.... and $\tilde{\mathcal{U}_1}\tilde{\mathcal{U}_0}\tilde{\mathcal{U}_1}$..... There are four possibilities for the beginning and ending of such a word and each one is mapped by the homomorphism to a different basis element (again using Proposition 3.8). It follows that $M = 0$. □

4. Translation of adelic to classical.

Let $\mathcal{G}$ be the set of all ordered pairs $(\alpha, \phi(z))$ where $\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathbb{R})^+$ and $\phi(z)$ is a holomorphic function on the upper half plane $\mathbb{H}$ such that $\phi(z)^2 = t \det(\alpha)^{-1/2}(cz + d)$ with t in the unit circle $S^1 := \{z \in \mathbb{C} : |z| = 1\}$. It is known [10] that $\mathcal{G}$ is a group under the following operation:

$$(\alpha, \phi(z)).(\beta, \psi(z)) = (\alpha\beta, \phi(\beta z)\psi(z)).$$

Let $P : \mathcal{G} \to \mathrm{GL}_2^+(\mathbb{R})$ be the homomorphism given by the projection map onto the first coordinate.

Let $k \geq 1$ and $\zeta = (\alpha, \phi(z))$. Define the slash operator $|[\zeta]_{k+1/2}$ on functions f on $\mathbb{H}$ by $f|[\zeta]_{k+1/2}(z) = f(\alpha z)(\phi(z))^{-2k-1}$.

Let N be divisible by 4 and $\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$. Define the automorphy factor

$$j(\alpha, z) = \varepsilon_d^{-1} \left(\frac{c}{d}\right) (cz + d)^{1/2},$$

where $\varepsilon_d = 1$ or $(-1)^{1/2}$ according as $d \equiv 1$ or $3 \pmod 4$ and $\left(\frac{c}{d}\right)$ is as in Shimura's notation. Let

$$\Delta_0(N) := \{\alpha^* = (\alpha, j(\alpha, z)) \in \mathcal{G} \mid \alpha \in \Gamma_0(N)\} \leq \mathcal{G}.$$

The map $L : \Gamma_0(N) \to \mathcal{G}$ given by $\alpha \mapsto \alpha^*$ defines an isomorphism onto $\Delta_0(N)$. Thus $P|_{\Delta_0(N)}$ and L are inverse of each other. Denote by $\Delta_0(N)$ and $\Delta_1(N)$ respectively the images of $\Gamma_0(N)$ and $\Gamma_1(N)$.

Let χ be an even Dirichlet character modulo N. Let $S_{k+1/2}(\Gamma_0(N), \chi)$ be the space of cusp forms of weight $k+1/2$, level N and character χ consisting of $f \in S_{k+1/2}(\Delta_1(N))$ such that $f|[\alpha^*]_{k+1/2}(z) = \chi(d)f(z)$ for all $\alpha \in \Gamma_0(N)$. In particular when χ is trivial $S_{k+1/2}(\Gamma_0(N), \chi) = S_{k+1/2}(\Delta_0(N))$.

Let ξ be an element of G such that $\Delta_0(N)$ and $\xi^{-1}\Delta_0(N)\xi$ are commensurable. We have an operator $|[\Delta_0(N)\xi\Delta_0(N)]_{k/2}$ on $S_{k+1/2}(\Delta_0(N))$ defined by

$$f|[\Delta_0(N)\xi\Delta_0(N)]_{k+1/2} = \det(\xi)^{(2k-3)/4} \sum_v f|[\xi_v]_{k+1/2}$$

where $\Delta_0(N)\xi\Delta_0(N) = \bigcup_v \Delta_0(N)\xi_v$.

Let $\mathbb{A} = \mathbb{A}_{\mathbb{Q}}$ be the adele ring of $\mathbb{Q}$ and $\widetilde{\mathrm{SL}_2}(\mathbb{A}) = \mathrm{SL}_2(\mathbb{A}) \times \{\pm 1\}$ with the group law: for $g = (g_\nu),\ h = (h_\nu) \in \mathrm{SL}_2(\mathbb{A})$ and $\epsilon_1,\ \epsilon_2 \in \{\pm 1\}$

$$(g, \epsilon_1)(h, \epsilon_2) = (gh,\ \epsilon_1\epsilon_2\sigma(g, h)),\ \text{where } \sigma(g, h) = \prod_\nu \sigma_\nu(g_\nu, h_\nu).$$

The group $\widetilde{\mathrm{SL}_2}(\mathbb{A})$ splits over $\mathrm{SL}_2(\mathbb{Q})$ and the splitting is given by

$$s_{\mathbb{Q}} : \mathrm{SL}_2(\mathbb{Q}) \longrightarrow \widetilde{\mathrm{SL}_2}(\mathbb{A}),\ g \mapsto (g, s_{\mathbb{A}}(g)) \text{ where } s_{\mathbb{A}}(g) = \prod_\nu s_\nu(g).$$

By [3, Proposition 2.16], for $\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_1(N)$, $s_{\mathbb{A}}(\alpha) = \left(\frac{c}{d}\right)_s$ unless $c = 0$ in which case $s_{\mathbb{A}}(\alpha) = 1$. Here $\left(\frac{c}{d}\right)_s = \left(\frac{c}{d}\right)(c, d)_\infty$.

Lemma 4.1. *Let $4 \mid N$. For $\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$, we have*

$$s_{\mathbb{A}}(\alpha) = \begin{cases} \left(\frac{c}{d}\right)_s (c, d)_2 & \text{if } c \neq 0 \text{ and } \mathrm{ord}_2(c) \text{ is even} \\ \left(\frac{c}{d}\right)_s & \text{if } c \neq 0 \text{ and } \mathrm{ord}_2(c) \text{ is odd} \\ 1 & \text{if } c = 0. \end{cases}$$

Proof. If $c = 0$ then $s_\nu(\alpha) = 1$ for all places ν and so $s_{\mathbb{A}}(\alpha) = 1$.

Suppose $c \neq 0$. Since $\alpha \in \Gamma_0(N)$ and $4 \mid N$, d is odd and coprime to c. By definition, for any finite prime q, we have $s_q(\alpha) = (c, d)_q$ if $\mathrm{ord}_q(c)$ is odd

and is 1 else. Hence

$$s_{\mathbb{A}}(\alpha) = \prod_{q \text{ finite}} s_q(\alpha) = \prod_{\mathrm{ord}_q(c) \text{ odd}} (c,d)_q .$$

It follows from the proof of [3, Proposition 2.16] (the proof only uses that d is odd and coprime to c), that $\left(\frac{c}{d}\right)_s = \prod_{q|c} (c,d)_q$. Now

$$\prod_{\mathrm{ord}_q(c) \text{ odd}} (c,d)_q = \prod_{q|c} (c,d)_q \prod_{\mathrm{ord}_q(c) \text{ even}>0} (c,d)_q = \left(\frac{c}{d}\right)_s \prod_{\mathrm{ord}_q(c) \text{ even}>0} (c,d)_q$$

So we just need to show that $\prod_{\mathrm{ord}_q(c) \text{ even}>0} (c,d)_q$ is $(c,d)_2$ if $\mathrm{ord}_2(c)$ is even and is 1 if $\mathrm{ord}_2(c)$ is odd (note that $\mathrm{ord}_2(c) \geq 2$). Let p be any odd prime such that $\mathrm{ord}_p(c)$ is even and > 0. Let $c = p^{2n}u$ where u is unit in $\mathbb{Z}_p$. Then $(c,d)_p = (u,d)_p = 1$ as both $d,\ u$ are units in $\mathbb{Z}_p$. Hence we are done. □

For $\tilde{g} = (\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \epsilon) \in \widetilde{\mathrm{SL}}_2(\mathbb{R})$ and $z \in \mathbb{H}$, define

$$\tilde{g}(z) = \frac{az+b}{cz+d} \qquad \text{and} \qquad J(\tilde{g},z) = \epsilon(cz+d)^{1/2}.$$

By [3, Lemma 3.3], $J(\tilde{g},z)$ satisfies automorphy condition i.e.,

$$J(\tilde{g}\tilde{h},z) = J(\tilde{g},\tilde{h}z)J(\tilde{h},z).$$

Let $f \in S_{k+1/2}(\Gamma_0(N))$ and $\alpha \in \Gamma_0(N)$. Then considering $\overline{\alpha} = (\alpha, s_{\mathbb{A}}(\alpha)) \in \widetilde{\mathrm{SL}}_2(\mathbb{R})$, using above lemma we have,

$$\begin{aligned} f(\overline{\alpha}z) &= \left(\frac{c}{d}\right)(\varepsilon_d^{-1})^{2k+1}(cz+d)^{k+1/2}f(z) \\ &= \left(\frac{c}{d}\right)(\varepsilon_d^{-1})^{2k+1}s_{\mathbb{A}}(\alpha)J(\overline{\alpha},z)^{2k+1}f(z) \\ &= \begin{cases} (\varepsilon_d^{-1}J(\overline{\alpha},z))^{2k+1}f(z) & \text{if } c=0 \text{ or } c\neq 0 \text{ and } \mathrm{ord}_2(c) \text{ is odd} \\ (c,d)_2\,(\varepsilon_d^{-1}J(\overline{\alpha},z))^{2k+1}f(z) & \text{if } c\neq 0 \text{ and } \mathrm{ord}_2(c) \text{ is even.} \end{cases} \end{aligned}$$

For $\theta \in \mathbb{R}$, let $k(\theta) = \begin{pmatrix} cos\theta & sin\theta \\ -sin\theta & cos\theta \end{pmatrix}$. Define $\tilde{K}_\infty := \{\tilde{k}(\theta) : \theta \in (-2\pi, 2\pi]\}$ where

$$\tilde{k}(\theta) = \begin{cases} (k(\theta),1) & \text{if } -\pi < \theta \leq \pi, \\ (k(\theta),-1) & \text{if } -2\pi < \theta \leq -\pi \text{ or } \pi < \theta \leq 2\pi. \end{cases}$$

Then $\tilde{K}_\infty$ is a maximal compact subgroup of $\widetilde{\mathrm{SL}}_2(\mathbb{R})$ and $\tilde{k}(\theta) \mapsto e^{i\frac{2k+1}{2}\theta}$ is a genuine character of $\tilde{K}_\infty$. Let

$$K_1(N) = \prod_{q<\infty} \{\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}_q) \ : \ c \equiv 0, \text{ and } a,d \equiv 1 \pmod{N\mathbb{Z}_q}\}.$$

Recall the strong approximation for $\widetilde{\mathrm{SL}}_2(\mathbb{A})$: Suppose $\tilde{g} = (g,\epsilon) \in \widetilde{\mathrm{SL}}_2(\mathbb{A})$. Then

$$\tilde{g} = (\alpha, s_{\mathbb{A}}(\alpha))\tilde{g}_\infty(k_1,1),$$

where $(\alpha, s_{\mathbb{A}}(\alpha)) \in s_{\mathbb{Q}}(\mathrm{SL}_2(\mathbb{Q}))$, $k_1 \in K_1(N)$ and $g_\infty^{\sim} \in \widetilde{\mathrm{SL}}_2(\mathbb{R})$ determined up to left multiplication by elements in $s_{\mathbb{Q}}(\Gamma_1(N))$.

We follow the notation of Waldspurger [12]. Let χ be an even Dirichlet character modulo N. Write $\chi_0 = \chi\left(\frac{-1}{\cdot}\right)^k$. Define $\tilde{\gamma_2}$ on $\mathbb{Z}_2^\times$ as

$$\tilde{\gamma_2}(t) = \begin{cases} 1 & \text{if } t \equiv 1 \pmod{4\mathbb{Z}_2} \\ -i & \text{if } t \equiv 3 \pmod{4\mathbb{Z}_2}, \end{cases}$$

and for $k_0 = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K_0^2(4)$, define

$$\tilde{\epsilon}_2(k_0) = \begin{cases} \tilde{\gamma_2}(d)^{-1} (c,d)_2 \, s_2(k_0) & \text{if } c \neq 0 \\ \tilde{\gamma_2}(d) & \text{if } c = 0. \end{cases}$$

Let χ_0 also denote the idelic character (of $\mathbb{Q}^\times \backslash \mathbb{A}_{\mathbb{Q}}^\times$) corresponding to the Dirichlet character χ_0 (it will be clear from the context when we consider χ_0 to be idelic or Dirichlet character). Let $A_{k+1/2}(N, \chi_0)$ denote the set of functions $\Phi : \widetilde{\mathrm{SL}}_2(\mathbb{A}) \to \mathbb{C}$ satisfying the following properties:

(1) $\Phi(s_{\mathbb{Q}}(\alpha)\tilde{g}(k_1, 1)) = \Phi(\tilde{g})$ for all $k_1 \in \prod_{q \nmid N} \mathrm{SL}_2(\mathbb{Z}_q), \alpha \in \mathrm{SL}_2(\mathbb{Q}), \tilde{g} \in \widetilde{\mathrm{SL}}_2(\mathbb{A})$.
(2) Φ is genuine, i.e., $\Phi((I, \zeta)\tilde{g}) = \zeta\Phi(\tilde{g})$ for $\zeta \in \mu_2$.
(3) For odd primes p such that $p^n \| N$, $\Phi(\tilde{g}(k_0, 1)) = \chi_{0,p}(d)\Phi(\tilde{g})$ for all $k_0 = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K_0^p(p^n)$.
(4) If $2^n \| N$ $(n \geq 2)$, $\Phi(\tilde{g}(k_0, 1)) = \tilde{\epsilon}_2(k_0)\chi_{0,2}(d)\Phi(\tilde{g})$ for all $k_0 \in K_0^2(2^n)$.
(5) $\Phi(\tilde{g}\tilde{k}(\theta)) = e^{i\frac{2k+1}{2}\theta}\Phi(\tilde{g})$ for all $\tilde{k}(\theta) \in \tilde{K}_\infty$.
(6) Φ is smooth as a function of $\widetilde{\mathrm{SL}}_2(\mathbb{R})$ and satisfies the differential equation $\Delta\Phi = -[(2k+1)/4 \cdot (2k-3)/4]\Phi$ where Δ is the Casimir operator.
(7) Φ is square integrable, that is $\int_{s_{\mathbb{Q}}(\mathrm{SL}_2(\mathbb{Q}))\backslash \widetilde{\mathrm{SL}}_2(\mathbb{A})/\mu_2} |\Phi(\tilde{g})|^2 d\tilde{g} < \infty$.
(8) Φ is cuspidal, that is $\int_{N_{\mathbb{Q}}\backslash N_{\mathbb{A}}} \Phi\left(\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \tilde{g}\right) da = 0$ for all $\tilde{g} \in \widetilde{\mathrm{SL}}_2(\mathbb{A})$.

In the above $\chi_{0,p}$ is the p-component of idelic character χ_0.

By [12, Proposition 3] there exists an isomorphism between

$$A_{k+1/2}(N, \chi_0) \to S_{k+1/2}(\Gamma_0(N), \chi)$$

given by $\Phi \mapsto f_\Phi$ where for $z \in \mathbb{H}$,

$$f_\Phi(z) = \Phi(\tilde{g}_\infty) J(\tilde{g}_\infty, i)^{2k+1}$$

where $\tilde{g}_\infty \in \widetilde{\mathrm{SL}}_2(\mathbb{R})$ is such that $\tilde{g}_\infty(i) = z$. The inverse map is given by $f \mapsto \Phi_f$ where for $g \in \widetilde{\mathrm{SL}}_2(\mathbb{A})$ if $\tilde{g} = (\alpha, s_{\mathbb{A}}(\alpha))\tilde{g}_\infty(k_1, 1)$,

$$\Phi_f(\tilde{g}) = f(\tilde{g}_\infty(i)) J(\tilde{g}_\infty, i)^{-2k-1}.$$

This isomorphism induces a ring isomorphism of spaces of linear operators,

$$q : \mathrm{End}_{\mathbb{C}}(A_{k+1/2}(N, \chi_0)) \to \mathrm{End}_{\mathbb{C}}(S_{k+1/2}(\Gamma_0(N), \chi))$$

given by

$$q(\mathcal{T})(f) = f_{\mathcal{T}(\Phi_f)}.$$

4.1. $N = 4M$, M **odd and** $p \| M$**.** Let p be an odd prime and let $N = 4M$ with M odd such that p strictly divides M. We shall translate the elements $\mathcal{T}_1, \mathcal{U}_1$ and $\mathcal{U}_0$ in the p-adic Hecke algebra to classical operators on $S_{k+1/2}(\Gamma_0(4M), \chi)$. We will restrict ourselves to χ being the trivial character modulo $4M$. In this case $\chi_0 = \left(\frac{-1}{\cdot}\right)^k$ is of conductor 1 or 4 and so $\chi_{0,p}$ is trivial on $\mathbb{Z}_p^\times$ while $\chi_{0,2}$ acts by $\chi_0^{-1} = \chi_0$ on $\mathbb{Z}_2^\times$.

Let γ be character on $(\mathbb{Z}_p/p\mathbb{Z}_p)^\times$ induced by $\chi_{0,p}|\mathbb{Z}_p^\times$ (so in the current case γ is trivial). Then Iwahori Hecke algebra $H(\overline{K_0^p(p)}, \gamma)$ is a subalgebra of $\mathrm{End}_{\mathbb{C}}(A_{k+1/2}(N, \chi_0))$ via the following action: for $\mathcal{T} \in H(\overline{K_0^p(p)}, \gamma)$ and $\Phi \in A_{k+1/2}(N, \chi_0)$,

$$\mathcal{T}(\Phi)(\tilde{g}) = \int_{\widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)} \mathcal{T}(\tilde{x})\Phi(\tilde{g}\tilde{x})d\tilde{x}.$$

Proposition 4.2. *Let χ be the trivial character modulo $4M$ with M as above and γ be induced by $\chi_{0,p}$. Let $\mathcal{T}_1$, $\mathcal{U}_1$ and $\mathcal{U}_0 \in H(\overline{K_0^p(p)}, \gamma)$ and $f \in S_{k+1/2}(\Gamma_0(4M), \chi)$. Then,*

(1) $q(\mathcal{T}_1)(f)(z) = \left(\dfrac{-1}{p}\right)^k p^{-k-1/2} \displaystyle\sum_{s=0}^{p^2-1} f\left(\frac{z+s}{p^2}\right)$.

(2) $q(\mathcal{U}_1)(f)(z) = \overline{\varepsilon_p}\left(\dfrac{-1}{p}\right)\left(\dfrac{M/p}{p}\right)\displaystyle\sum_{s=0}^{p-1} f|[(\alpha_s, \phi_{\alpha_s})]_{k+1/2}(z)$, *where*

$\alpha_s = \begin{pmatrix} p^2n - 4Mms & m \\ 4pM(1-s) & p \end{pmatrix} \in \mathrm{M}_2(\mathbb{Z})$ *is of determinant* p^2 *and* $m,\ n \in \mathbb{Z}$ *are such that* $pn - (4M/p)m = 1$, *and* $\phi_{\alpha_s}(z) = (4M(1-s)z+1)^{1/2}$.

(3) $q(\mathcal{U}_0)(f)(z) = \displaystyle\sum_{s=0}^{p-1} f|[(\beta_s, \phi_{\beta_s})]_{k+1/2}(z)$, *where*

$\beta_s = \begin{pmatrix} 1 & m-s \\ 4M_1 & np - 4M_1 s \end{pmatrix} \in \Gamma_0(4M_1)$ *with* $M_1 = M/p$ *and* $m,\ n \in \mathbb{Z}$ *are chosen as above and* $\phi_{\beta_s} = (4M_1 z + (np - 4M_1 s))^{1/2}$.

Proof. For (1), let $\tilde{g}_\infty = (g_\infty, 1) \in \widetilde{\mathrm{SL}}_2(\mathbb{R})$ such that $\tilde{g}_\infty i = z$. Then using decomposition in Lemma 3.5 we have

$$\mathcal{T}_1(\Phi_f)(\tilde{g_\infty}) = \sum_{s=0}^{p^2-1} \overline{\gamma}(h(p), 1)\Phi_f(\tilde{g}_\infty(x(s), 1)(h(p), 1))$$

$$= \overline{\varepsilon_p} \sum_{s=0}^{p^2-1} \Phi_f(\tilde{g}_\infty(x(s),1)(h(p),1)).$$

Take $A_s = h(p^{-1})x(-s) = \begin{pmatrix} p^{-1} & -p^{-1}s \\ 0 & p \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Q})$, then $s_{\mathbb{Q}}(A_s) = (A_s, 1)$. Now the ∞-component of

$$\underbrace{(A_s,1)}_{\text{diagonal emb.}} \cdot \underbrace{\tilde{g}_\infty}_{\infty \text{ place}} \underbrace{\cdot(x(s),1)(h(p),1)}_{p \text{ place}}$$

is $(A_s,1)\tilde{g}_\infty$, for a prime q such that $(q,2M)=1$ the q-component is $(A_s,1) \in \mathrm{SL}_2(\mathbb{Z}_q)\times\{1\}$, for a prime r such that $(r,2p)=1$ and $r^b \| M$, the r-component is $(A_s,1) \in K_0^r(r^b)\times\{1\}$, the 2-component is $(A_s,1) \in K_0^2(4)\times\{1\}$ and the p-component is $(A_s,1)(x(s),1)(h(p),1) = (I,(p,p)_p) = (I, \left(\frac{-1}{p}\right))$.

Since χ is trivial, $\chi_{0,2} = \left(\frac{-1}{\cdot}\right)^k$ while $\chi_{0,p}$ and $\chi_{0,r}$ are trivial. So the 2-component acts by $\tilde{\epsilon}_2(A_2)\chi_{0,2}(p) = \tilde{\gamma_2}(p)\chi_{0,2}(p) = \overline{\varepsilon_p}\left(\frac{-1}{p}\right)^k$. Thus,

$$\begin{aligned}
\mathcal{T}_1(\Phi_f)(\tilde{g}_\infty) &= \overline{\varepsilon_p} \sum_{s=0}^{p^2-1} \Phi_f(s_{\mathbb{Q}}(A)\tilde{g}_\infty(x(s),1)(h(p),1)) \\
&= (\overline{\varepsilon_p})^2 \left(\frac{-1}{p}\right)^k \left(\frac{-1}{p}\right) \sum_{s=0}^{p^2-1} \Phi_f(Ag_\infty, 1) \\
&= \left(\frac{-1}{p}\right)^k \sum_{s=0}^{p^2-1} f(Ag_\infty(i)) J((Ag_\infty,1),i)^{-2k-1}.
\end{aligned}$$

Consequently,

$$q(\mathcal{T}_1)(f)(z) = \mathcal{T}_1(\Phi_f)(\tilde{g}_\infty) J((g_\infty,1),i)^{2k+1} = \left(\frac{-1}{p}\right)^k p^{-k-1/2} \sum_{s=0}^{p^2-1} f(\frac{z+s}{p^2}).$$

For (2) we need the following decomposition (we use $(4,M)=1$)

$$K_0 w(p^{-1}) K_0 = \bigcup_{s \in \mathbb{Z}_p/p\mathbb{Z}_p} y(4Ms) w(p^{-1}) K_0$$

Taking $\tilde{g}_\infty$ such that $\tilde{g}_\infty i = z$ we have

$$\mathcal{U}_1(\Phi_f)(\tilde{g}_\infty) = \overline{\varepsilon_p}\left(\frac{-1}{p}\right) \sum_{s=0}^{p-1} \Phi_f(\tilde{g}_\infty(y(4Ms),1)(w(p^{-1}),1)).$$

Since p is coprime to $4M/p$, we fix $m,\ n \in \mathbb{Z}$ such that $pn - (4M/p)m = 1$. For $0 \le s \le p-1$, take

$$A_s = \begin{pmatrix} pn & \frac{m}{p} \\ 4M & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -4Ms & 1 \end{pmatrix} = \begin{pmatrix} pn - 4ms\frac{M}{p} & \frac{m}{p} \\ 4M(1-s) & 1 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Q}).$$

Since $s_\nu(A_s)=1$ for all primes ν we have $s_{\mathbb{Q}}(A_s)=(A_s,1)$. As before the ∞-component of

$$s_{\mathbb{Q}}(A_s)\ \tilde{g}_\infty\ (y(4Ms),1)(w(p^{-1}),1)$$

is $(A_s,1)\tilde{g}_\infty$, for a prime q such that $(q,2M)=1$ the q-component is $(A_s,1)\in \mathrm{SL}_2(\mathbb{Z}_q)\times\{1\}$, for a prime r such that $(r,2p)=1$ and $r^b\|M$, the r-component is $(A_s,1)\in K_0^r(r^b)\times\{1\}$, the 2-component is $(A_s,1)\in K_1^2(4)\times\{1\}$ (as $(2,2)$-th entry of A_s is 1). At the p-component we check that $(A_s,1)=(\begin{pmatrix} pn & m/p \\ 4M & 1\end{pmatrix},1)(y(-4Ms),1)$ and

$$(A_s,1)(y(4Ms),1)(w(p^{-1}),1)=(\begin{pmatrix} -m & n \\ -p & 4M/p\end{pmatrix},\left(\frac{M/p}{p}\right)).$$

Since χ is trivial, the p and r components acts trivially and the 2-component acts by $\tilde{\epsilon}_2(A_s)\chi_{0,2}(1)=1$. Hence

$$\begin{aligned}\mathcal{U}_1(\Phi_f)(\tilde{g}_\infty)&=\overline{\varepsilon_p}\left(\frac{-1}{p}\right)\sum_{s=0}^{p-1}\Phi_f(s_{\mathbb{Q}}(A_s)\tilde{g}_\infty(y(4ps),1)(w(p^{-1}),1))\\&=\overline{\varepsilon_p}\left(\frac{-1}{p}\right)\left(\frac{M/p}{p}\right)\sum_{s=0}^{p-1}\Phi_f((A_s,1)(g_\infty,1))\end{aligned}$$

$$=\overline{\varepsilon_p}\left(\frac{-1}{p}\right)\left(\frac{M/p}{p}\right)\sum_{s=0}^{p-1}f((A_s,1)z)J((A_s,1),z)^{-2k-1}J((g_\infty,1),i)^{-2k-1}.$$

So we have

$$q(\mathcal{U}_1)(f)(z)=\overline{\varepsilon_p}\left(\frac{-1}{p}\right)\left(\frac{M/p}{p}\right)\sum_{s=0}^{p-1}f((A_s,1)z)J((A_s,1),z)^{-2k-1}.$$

Let $\alpha_s=A_s\begin{pmatrix} p & 0\\ 0 & p\end{pmatrix}$ and $\phi_{\alpha_s}(z)=(4M(1-s)z+1)^{1/2}$. Then $q(\mathcal{U}_1)(f)(z)=$

$$=\overline{\varepsilon_p}\left(\frac{-1}{p}\right)\left(\frac{M/p}{p}\right)\sum_{s=0}^{p-1}f\left(\frac{(p^2n-4mMs)z+m}{4pM(1-s)z+p}\right)(4M(1-s)z+1)^{-k-1/2}$$

$$=\overline{\varepsilon_p}\left(\frac{-1}{p}\right)\left(\frac{M/p}{p}\right)\sum_{s=0}^{p-1}f|[(\alpha_s,\phi_{\alpha_s})]_{k+1/2}(z).$$

For (3), using Lemma 3.5 we have

$$\mathcal{U}_0(\Phi_f)(\tilde{g}_\infty)=\sum_{s=0}^{p-1}\Phi_f(\tilde{g}_\infty(x(s),1)(w(1),1)).$$

Let $m,\ n\in\mathbb{Z}$ such that $pn-(4M/p)m=1$ and let $M_1=M/p$. For $0\le s\le p-1$, take

$$A_s=\begin{pmatrix} 1 & -s+m\\ 4M_1 & -4M_1s+np\end{pmatrix}\in\Gamma_0(4M_1).$$

Since the Kronecker symbol $\left(\frac{4M_1}{-4M_1s+np}\right) = \left(\frac{M_1}{-4M_1s+np}\right) = \left(\frac{-4M_1s+np}{M_1}\right) = \left(\frac{np}{M_1}\right) = 1$, by Lemma 4.1 we get that

$$s_{\mathbb{A}}(A_s) = \begin{cases} (4M_1, -4M_1s+np)_2 & \text{if } \mathrm{ord}_2(M_1) \text{ even} \\ 1 & \text{if } \mathrm{ord}_2(M_1) \text{ odd.} \end{cases}$$

The ∞-component of $s_{\mathbb{Q}}(A_s)$ $\tilde{g}_\infty$ $(x(s),1)(w(1),1)$ is $(A_s, s_{\mathbb{A}}(A_s))(g_\infty, 1)$, for a prime q such that $(q, 2M) = 1$ the q-component is $(A_s, 1) \in \mathrm{SL}_2(\mathbb{Z}_q) \times \{1\}$, if r is an odd prime such that $r^b \| M$ then the r-component is $(A_s, 1) \in K_0^r(r^b) \times \{1\}$ and the 2-component is $(A_s, 1) \in K_1^2(4) \times \{1\}$. For p-component, since $\mathrm{ord}_p(4M_1) = 0$, we have $(\begin{pmatrix} 1 & m \\ 4M_1 & np \end{pmatrix}, 1)(x(-s), 1) = (A_s, 1)$ and $(\begin{pmatrix} 1 & m \\ 4M_1 & np \end{pmatrix}, 1)(w(1), 1) = (\begin{pmatrix} -m & 1 \\ -np & 4M_1 \end{pmatrix}, \beta)$ where β is either $(4M_1, -1)_p$ or $(4M_1, np)_p$ depending on whether $\mathrm{ord}_p(np)$ is odd or even. In either case it is clear that β is 1. Thus the p-component is $(\begin{pmatrix} -m & 1 \\ -np & 4M_1 \end{pmatrix}, 1) \in K_0 \times \{1\}$

Since χ is trivial, the p and r components acts trivially, and the 2-component acts by $\tilde{\epsilon}_2(A_s)\chi_{0,2}(-4M_1s+np) = (4M_1, -4M_1s+np)_2 s_2(A_s)$ which is precisely equal to $s_{\mathbb{A}}(A_s)$.

Thus $\mathcal{U}_0(\Phi_f)(\tilde{g}_\infty) =$

$$= \sum_{s=0}^{p-1} \Phi_f((A_s,1)\tilde{g}_\infty) = \sum_{s=0}^{p-1} f(A_s z) J((A_s,1),z)^{-2k-1} J((g_\infty,1),i)^{-2k-1}$$

and consequently

$$q(\mathcal{U}_0)(f)(z) = \sum_{s=0}^{p-1} f\left(\frac{z+(m-s)}{4M_1z+(np-4M_1s)}\right)(4M_1z+(np-4M_1s))^{-k-1/2}.$$

□

Recall by [10, Proposition 1.5], for $m \mid M$, the map U_m given by

$$U_m\left(\sum_{n=1}^{\infty} a_n q^n\right) = \sum_{n=1}^{\infty} a_{mn} q^n = m^{(2k-3)/4} \sum_{s=0}^{m-1} f|[(\begin{pmatrix} 1 & s \\ 0 & m \end{pmatrix}, m^{1/4})]_{k+1/2}(z)$$

sends $S_{k+1/2}(\Gamma_0(4M))$ to $S_{k+1/2}(\Gamma_0(4M), (\frac{m}{\cdot}))$. Thus by (1) of the above proposition, we have $q(\mathcal{T}_1)(f)(z) = \left(\frac{-1}{p}\right)^k p^{(3-2k)/2} U_{p^2}(f)$.

Let us denote $q(p^{-1/2}\mathcal{U}_1)$ by $\widetilde{W}_{p^2}$ and $q(\mathcal{U}_0)$ by $\widetilde{Q}_p$.

Corollary 4.3. *On $S_{k+1/2}(\Gamma_0(4M))$ we have*

(1) $\widetilde{W}_{p^2}$ is an involution.
(2) $(\widetilde{Q}_p - p)(\widetilde{Q}_p + 1) = 0$.
(3) $\widetilde{Q}_p = \left(\frac{-1}{p}\right)^k p^{1-k} U_{p^2} \widetilde{W}_{p^2}$.

(4) If $f \in S_{k+1/2}(\Gamma_0(4M/p))$ then $\widetilde{Q}_p(f) = pf$.

Proof. The proof of (1)-(4) follows by using Proposition 3.9 and 4.2. For (5) we use Proposition 4.2(3). □

We further define an operator $\widetilde{Q}'_p$ on $S_{k+1/2}(\Gamma_0(4M))$ to be the conjugate of $\widetilde{Q}_p$ by $\widetilde{W}_{p^2}$, i.e., $\widetilde{Q}'_p = \widetilde{W}_{p^2}\widetilde{Q}_p\widetilde{W}_{p^2}$. Thus $\widetilde{Q}'_p$ satisfies same quadratic as $\widetilde{Q}_p$ and we have $\widetilde{Q}'_p = \left(\frac{-1}{p}\right)^k p^{1-k}\widetilde{W}_{p^2}U_{p^2}$.

Remark 3. *We note that for a prime q such that $(q, 2M) = 1$, one can similarly obtain the usual Hecke operator T_{q^2} on $S_{k+1/2}(\Gamma_0(4M))$. In particular, if we take $\mathcal{T}_1 := X_{(h(q),1)} \in H(\overline{\mathrm{SL}_2(\mathbb{Z}_q)}, \gamma_q)$ then $q(\mathcal{T}_1) = \left(\frac{-1}{p}\right)^k p^{(3-2k)/2}T_{q^2}$.*

Moreover if p and q are distinct primes such that p^n, q^m strictly divides N then the operators $\mathcal{S} \in H(\overline{K_0^p(p^n)}, \gamma_p)$ and $\mathcal{T} \in H(\overline{K_0^q(q^m)}, \gamma_q)$ in $\mathrm{End}_{\mathbb{C}}(S_{k+1/2}(\Gamma_0(N)))$ commute.

In particular the operators $\widetilde{Q}_p$, $\widetilde{W}_{p^2}$ on $S_{k+1/2}(\Gamma_0(4M))$ that we defined above commute with T_{q^2} for primes q coprime to $2M$.

4.2. **self-adjointness.** In this subsection we check that the operators $\widetilde{W}_{p^2}$, $\widetilde{Q}_p$ and $\widetilde{Q}'_p$ are self-adjoint operators on $S_{k+1/2}(\Gamma_0(4M))$.

Proposition 4.4. *The operator $\widetilde{W}_{p^2}$ is self-adjoint with respect to Petersson inner product.*

Proof. We write

$$\widetilde{W}_{p^2}(f) = \frac{\overline{\varepsilon_p}}{\sqrt{p}}\left(\frac{-1}{p}\right)\left(\frac{M/p}{p}\right)\mathcal{S}_p(f), \quad \mathcal{S}_p(f) := \sum_{s\in\mathbb{Z}/p\mathbb{Z}} f|[(\alpha_s, \phi_{\alpha_s}(z))]_{k+1/2}$$

where $(\alpha_s, \phi_{\alpha_s}(z)) = \left(\begin{pmatrix} p^2n - 4mMs & m \\ 4pM(1-s) & p\end{pmatrix}, (4M(1-s)z+1)^{1/2}\right) \in \mathcal{G}$ and $n,\ m$ are integers such that $pn - (4M/p)m = 1$.

We will show that $\langle \mathcal{S}_p(f), g\rangle = \left(\frac{-1}{p}\right)\langle f, \mathcal{S}_p(g)\rangle$. We write $\mathcal{S}_p = S_{1,p} + S_{2,p}$ where $S_{1,p}$ consists of $s = 0$ term and $S_{2,p}$ consists of rest of the terms. Also let $M_1 = M/p$.

We first consider $S_{2,p}$. For $s \neq 0$, as $pn - 4M_1ms = 1 + 4M_1m(1-s)$ it is clear that $pn - 4M_1ms$ and $4M(1-s)$ are relatively coprime, hence there exists integers u, v such that $u(-pn + 4M_1ms) - v4M(1-s) = 1$. Let $X = \begin{pmatrix} u & v \\ 4M(1-s) & -pn + 4M_1ms\end{pmatrix} \in \Gamma_0(4M)$, then $X^* = (X, j(X,z))$ where $j(X,z) = (-1)^{-1/2}\left(\frac{4M(1-s)}{-pn+4M_1ms}\right)(4M(1-s)z + (-pn+4M_1ms))^{1/2}$ as $-pn + 4M_1ms \equiv -1 \pmod 4$. Since f has level $4M$ we have $f|[(\alpha_s, \phi_{\alpha_s}(z))]_{k+1/2} = f|[X^*]_{k+1/2}|[(\alpha_s, \phi_{\alpha_s}(z))]_{k+1/2}$. We claim that in $\mathcal{G}$,

$$X^* \cdot (\alpha_s, \phi_{\alpha_s}(z)) = \left(\begin{pmatrix} -p & um + vp \\ 0 & -p\end{pmatrix}, -\left(\frac{um}{p}\right)\left(\frac{M_1}{p}\right)\right).$$

It is easy to see equality in the matrix component, so just need to check that $j(X,\alpha_s z)\cdot\phi_{\alpha_s}(z) = -\left(\frac{um}{p}\right)\left(\frac{M_1}{p}\right)$. We see that $j(X,\alpha_s z)$ simplifies to $(-1)^{-1/2}\left(\frac{4M(1-s)}{-pn+4M_1ms}\right)\left(\frac{-1}{4M(1-s)z+1}\right)^{1/2}$ and so $j(X,\alpha_s z)\cdot\phi_{\alpha_s}(z) = \left(\frac{4M(1-s)}{-pn+4M_1ms}\right)$. Thus we are left to show equality of the Kronecker symbols $\left(\frac{4M(1-s)}{-pn+4M_1ms}\right) = -\left(\frac{um}{p}\right)\left(\frac{M_1}{p}\right)$. Note that $\left(\frac{m}{p}\right)\left(\frac{M_1}{p}\right) = \left(\frac{-1}{p}\right)$ and $\left(\frac{u}{p}\right) = \left(\frac{-1-4M_1m(1-s)}{p}\right)$, so we have $\left(\frac{um}{p}\right)\left(\frac{M_1}{p}\right) = \left(\frac{p}{-1-4M_1m(1-s)}\right)$. Further $\left(\frac{4M(1-s)}{-pn+4M_1ms}\right) = \left(\frac{p}{-1-4M_1m(1-s)}\right)\left(\frac{-4M_1m(1-s)}{-1-4M_1m(1-s)}\right)\left(\frac{-m}{-1-4M_1m(1-s)}\right) = \left(\frac{p}{-1-4M_1m(1-s)}\right)\left(\frac{-m}{-1-4M_1m(1-s)}\right)$. We now check that $\left(\frac{-m}{-1-4M_1m(1-s)}\right) = -1$. Suppose m is odd, then $\left(\frac{-m}{-1-4mM_1(1-s)}\right) = -\left(\frac{m}{-1-4M_1m(1-s)}\right) = -\left(\frac{-1-4M_1m(1-s)}{m}\right)\left(\frac{-1}{m}\right) = -1$. Suppose $m = 2^km'$ where $k>1$ and m' odd, then

$$\left(\frac{-m}{-1-4M_1m(1-s)}\right) = \begin{cases}\left(\frac{-m'}{-1-4M_12^km'(1-s)}\right) & \text{if } k \text{ even}\\ \left(\frac{-2m'}{-1-4M_12^km'(1-s)}\right) & \text{if } k \text{ odd}\end{cases} = \left(\frac{-m'}{-1-4M_12^km'(1-s)}\right) = -1$$

(as $\left(\frac{2}{r}\right) = (-1)^{(r^2-1)/8}$). Thus our claim is proved.

Hence $f|[(\alpha_s,\phi_{\alpha_s}(z))]_{k+1/2} = f|[(\begin{pmatrix}-p & um\\ 0 & -p\end{pmatrix}, -\left(\frac{um}{p}\right)\left(\frac{M_1}{p}\right))]_{k+1/2}$ and consequently

$$S_{2,p}(f) = -\left(\frac{M_1}{p}\right)\sum_{u\in(\mathbb{Z}/p\mathbb{Z})^\times} f|[(\begin{pmatrix}-p & u\\ 0 & -p\end{pmatrix}, \left(\frac{u}{p}\right))]_{k+1/2}.$$

Since the adjoint of $|[(\begin{pmatrix}-p & u\\ 0 & -p\end{pmatrix}, \left(\frac{u}{p}\right))]_{k+1/2}$ is $|[(\begin{pmatrix}-p & -u\\ 0 & -p\end{pmatrix}, \left(\frac{u}{p}\right))]_{k+1/2}$, so the adjoint of $S_{2,p}(f)$ is $\left(\frac{-1}{p}\right)S_{2,p}(f)$, i.e., $\langle S_{2,p}(f),g\rangle = \left(\frac{-1}{p}\right)\langle f, S_{2,p}(g)\rangle$.

Next we consider the term $S_{1,p}(f) = f|[(\begin{pmatrix}p^2n & m\\ 4pM & p\end{pmatrix}, (4Mz+1)^{1/2})]_{k+1/2}$. Let $\gamma_p := \begin{pmatrix}a & b\\ c & d\end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ such that $\gamma_p \equiv \begin{pmatrix}0 & -1\\ 1 & 0\end{pmatrix} \pmod p$, $\equiv \begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix} \pmod 8$. We claim that

$$S_{1,p}(f) = f|[(\begin{pmatrix}pa & b\\ p^2c & pd\end{pmatrix}, \left(\frac{M_1}{p}\right)\left(\frac{c}{d}\right)(cpz+d)^{1/2})]_{k+1/2}$$

Choose $Y = \begin{pmatrix}a-4bM_1 & \frac{-ma+bpn}{p}\\ pc-4Md & -mc+dpn\end{pmatrix} \in \Gamma_0(4M)$. To prove the claim we need to check that

$$Y^*\cdot(\begin{pmatrix}p^2n & m\\ 4pM & p\end{pmatrix}, (4Mz+1)^{1/2}) = (\begin{pmatrix}pa & b\\ p^2c & pd\end{pmatrix}, \left(\frac{M_1}{p}\right)\left(\frac{c}{d}\right)(cpz+d)^{1/2}).$$

As before, matrix equality is easy to check and the automorphy factor of the left hand side equals kronecker symbol $\left(\frac{pc-4Md}{-cm+dpn}\right)$ times $(pcz+d)^{1/2}$. So we need to show that $\left(\frac{pc-4Md}{-cm+dpn}\right) = \left(\frac{M_1}{p}\right)\left(\frac{c}{d}\right)$. Since $d-m(c-4M_1d) = -cm+dpn \equiv 1 \pmod 4$ we have $\left(\frac{pc-4Md}{-cm+dpn}\right) = \left(\frac{-cm+dpn}{p}\right)\left(\frac{c-4M_1d}{-cm+dpn}\right) = \left(\frac{-mc}{p}\right)\left(\frac{-m(c-4M_1d)}{d-m(c-4M_1d)}\right)\left(\frac{-m}{-cm+dpn}\right) = \left(\frac{-m}{p}\right)\left(\frac{-d}{d-m(c-4M_1d)}\right)\left(\frac{m}{d-m(c-4M_1d)}\right) = \left(\frac{M_1}{p}\right)\left(\frac{d-m(c-4M_1d)}{d}\right)\left(\frac{m}{d-m(c-4M_1d)}\right) = \left(\frac{M_1}{p}\right)\left(\frac{c}{d}\right)\left(\frac{m}{d}\right)\left(\frac{m}{d-m(c-4d)}\right)$. We can check $\left(\frac{m}{d}\right) = \left(\frac{m}{d-m(c-4d)}\right)$ as before by considering m odd and even case (when m is even we use that $\left(\frac{2}{d}\right)=1$). Thus our claim is proved.

Now note that

$$\left(\begin{pmatrix} pa & b \\ p^2c & pd\end{pmatrix}, \left(\frac{c}{d}\right)(cpz+d)^{1/2}\right) = \left(\begin{pmatrix}1&0\\0&p\end{pmatrix}, p^{1/4}\right)\cdot\gamma_p^*\cdot\left(\begin{pmatrix}p&0\\0&1\end{pmatrix}, p^{-1/4}\right) =: \varsigma_p,$$

and so $S_{1,p}(f) = \left(\frac{M_1}{p}\right) f|[\varsigma_p]_{k+1/2}$.

We check similarly that

$$f|[\left(\begin{pmatrix}1&0\\0&p\end{pmatrix}, p^{1/4}\right)\cdot(\gamma_p^*)^2\cdot\left(\begin{pmatrix}p&0\\0&1\end{pmatrix}, p^{-1/4}\right)]_{k+1/2} = \left(\frac{-1}{p}\right) f.$$

By the above action on f we get that $f|[\varsigma_p^{-1}]_{k+1/2} = \left(\frac{-1}{p}\right) f|[\varsigma_p]_{k+1/2}$. Since the adjoint of ς_p is ς_p^{-1} we get $\langle S_{1,p}(f), g\rangle = \left(\frac{-1}{p}\right)\langle f, S_{1,p}(g)\rangle$.

Thus $\langle \mathcal{S}_p(f), g\rangle = \left(\frac{-1}{p}\right)\langle f, \mathcal{S}_p(g)\rangle$. So $\langle \widetilde{W}_{p^2}(f), g\rangle = \frac{\overline{\varepsilon_p}}{\sqrt{p}}\left(\frac{-M_1}{p}\right)\langle \mathcal{S}_p(f), g\rangle$ $= \frac{\overline{\varepsilon_p}}{\sqrt{p}}\left(\frac{M_1}{p}\right)\langle f, \mathcal{S}_p(g)\rangle = \langle f, \frac{\varepsilon_p}{\sqrt{p}}\left(\frac{M_1}{p}\right)\mathcal{S}_p(g)\rangle = \langle f, \widetilde{W}_{p^2}(g)\rangle$. Hence we are done. □

Remark 4. *Recall our earlier remark 2 where we checked that our p-adic operator $\mathcal{U}_1' = \overline{\varepsilon_p}\mathcal{U}_1$ and Ueda's classical operator Y_p satisfy same relations. Since $q(\mathcal{U}_1') = \mathcal{S}_p$, it is natural to compare $\mathcal{S}_p$ and Y_p. While $\mathcal{S}_p = S_{1,p} + S_{2,p}$, it turns out from the above proof that Ueda's Y_p becomes equal to $\left(\frac{M/p}{p}\right)(S_{1,p} - S_{2,p})$.*

Next we want to show that $\widetilde{Q}_p = q(\mathcal{U}_0)$ is self-adjoint. We use the relations $\mathcal{U}_1\mathcal{T}_1\mathcal{U}_1 = p\mathcal{T}_{-1}$ and $\mathcal{T}_1\mathcal{U}_1 = p\,\mathcal{U}_0$. So we have

$$\langle q(\mathcal{U}_0)f, g\rangle = \frac{1}{p}\,\langle q(\mathcal{T}_1)q(\mathcal{U}_1)f, g\rangle.$$

Since by above theorem $q(\mathcal{U}_1)$ is self-adjoint we have

$$\langle f, q(\mathcal{U}_0)g\rangle = \frac{1}{p}\,\langle f, p\; q(\mathcal{U}_0)g\rangle = \frac{1}{p}\,\langle f, q(\mathcal{T}_1)q(\mathcal{U}_1)g\rangle$$

$$= \frac{1}{p}\,\langle f, \frac{1}{p}\; q(\mathcal{U}_1)^2q(\mathcal{T}_1)q(\mathcal{U}_1)g\rangle = \frac{1}{p}\,\langle q(\mathcal{U}_1)f, \frac{1}{p}\; q(\mathcal{U}_1)q(\mathcal{T}_1)q(\mathcal{U}_1)g\rangle$$

$$= \frac{1}{p}\ \langle q(\mathcal{U}_1)f, q(\mathcal{T}_{-1})g\rangle.$$

Since $q(\mathcal{U}_1)$ is surjective we get that $q(\mathcal{U}_0)$ is self-adjoint *iff* $\langle q(\mathcal{T}_1)q(\mathcal{U}_1)f, g\rangle = \langle q(\mathcal{U}_1)f, q(\mathcal{T}_{-1})g\rangle$, i.e., *iff* the adjoint of $q(\mathcal{T}_{-1})$ is $q(\mathcal{T}_1)$.

In order to obtain the adjoint of $q(\mathcal{T}_{-1})$ we first compute its action on $S_{k+1/2}(\Gamma_0(4M))$.

Proposition 4.5. *Let χ be the trivial character and γ induced by $\chi_{0,p}$. For $\mathcal{T}_{-1} = X_{(h(p^{-1}),1)} \in H(\overline{K_0^p(p)}, \gamma)$ and $f \in S_{k+1/2}(\Gamma_0(4M), \chi)$, we have*

$$q(\mathcal{T}_{-1})(f)(z) = \left(\frac{-1}{p}\right)^k \sum_{s=0}^{p^2-1} f|[(\alpha_s, \phi_{\alpha_s}(z))]_{k+1/2}(z)$$

where $\alpha_s = \begin{pmatrix} p^2 & 0 \\ -4Ms & 1 \end{pmatrix}$ *and* $\phi_{\alpha_s}(z) = (-4(M/p)sz + p^{-1})^{1/2}$.

Proof. Let $\tilde{g}_\infty = (g_\infty, 1) \in \widetilde{\mathrm{SL}_2}(\mathbb{R})$ such that $\tilde{g}_\infty i = z$. Then using the decomposition $K_0 h(p^{-1}) K_0 = \bigcup_{s\in\mathbb{Z}_p/p^2\mathbb{Z}_p} y(4Ms)h(p^{-1})K_0$ we have

$$\mathcal{T}_{-1}(\Phi_f)(\tilde{g_\infty}) = \sum_{s=0}^{p^2-1} \overline{\gamma}(h(p^{-1}),1)\Phi_f(\tilde{g}_\infty(y(4Ms),1)(h(p^{-1}),1))$$

$$= \overline{\varepsilon_p} \sum_{s=0}^{p^2-1} \Phi_f(\tilde{g}_\infty(y(4Ms),1)(h(p^{-1}),1)).$$

Take $A_s = h(p)y(-4Ms) = \begin{pmatrix} p & 0 \\ -4(M/p)s & p^{-1} \end{pmatrix}$, then $s_{\mathbb{Q}}(A_s) = (A_s, \xi_s)$ where $\xi_s := \begin{cases} 1 & \text{if } s = 0 \\ 1 & \text{if } \mathrm{ord}_p(s) = 1 \text{ and } \mathrm{ord}_2(s) \text{ odd} \\ \left(\frac{-1}{p}\right)(\frac{Ms}{p}, p)_2 & \text{if } \mathrm{ord}_p(s) = 1 \text{ and } \mathrm{ord}_2(s) \text{ even} \\ \left(\frac{-1}{p}\right)(\frac{Ms}{p}, p)_p & \text{if } (s,p) = 1 \text{ and } \mathrm{ord}_2(s) \text{ odd} \\ (\frac{Ms}{p}, p)_2\ (\frac{Ms}{p}, p)_p & \text{if } (s,p) = 1 \text{ and } \mathrm{ord}_2(s) \text{ even.} \end{cases}$

Thus

$$\mathcal{T}_{-1}(\Phi_f)(\tilde{g_\infty}) = \overline{\varepsilon_p} \sum_{s=0}^{p^2-1} \xi_s \Phi_f((A_s,1)\tilde{g}_\infty(y(4Ms),1)(h(p^{-1}),1)).$$

Now the ∞-component of $(A_s,1)\ \tilde{g}_\infty\ (y(4Ms),1)(h(p^{-1}),1)$ is $(A_s,1)\tilde{g}_\infty$, for a prime q such that $(q,2M) = 1$ the q-component is $(A_s,1) \in \mathrm{SL}_2(\mathbb{Z}_q)\times\{1\}$, if r is an odd prime coprime to p such that $r^b \| M$ then the r-component belongs to $K_0^r(r^b) \times \{1\}$, the 2-component is $(\begin{pmatrix} p & 0 \\ -4(M/p)s & p^{-1} \end{pmatrix}, 1) \in K_0^2(4) \times \{1\}$

and the p-component is $(A_s,1)(y(4Ms),1)(h(p^{-1}),1)$ which is precisely equal to (I,η_s) where $\eta_s := \begin{cases} \left(\frac{-1}{p}\right) & \text{if } s=0 \\ 1 & \text{if } \mathrm{ord}_p(s)=1 \\ \left(\frac{-1}{p}\right)(\frac{Ms}{p},p)_p & \text{if } (s,p)=1. \end{cases}$

Since χ is trivial, $\chi_{0,p}$ is trivial and so the p-component acts on Φ_f simply by multiplication by η_s. Next we look at how the 2-component acts on Φ_f. Since $\chi_{0,2}=\left(\frac{-1}{\cdot}\right)^k$ we get that

$$\tilde{\epsilon}_2(A_s)\chi_{0,2}(p^{-1}) = \begin{cases} \tilde{\gamma_2}(p^{-1})\chi_{0,2}(p^{-1}) & \text{if } s=0 \\ \tilde{\gamma_2}(p^{-1})^{-1}(-4\frac{M}{p}s,p^{-1})_2 s_2(A_s)\chi_{0,2}(p^{-1}) & \text{if } s\neq 0 \end{cases}$$

$$= \begin{cases} \overline{\varepsilon_p}\left(\frac{-1}{p}\right)^k & \text{if } s=0 \\ \varepsilon_p\left(\frac{-1}{p}\right)^k & \text{if } s\neq 0 \text{ and } \mathrm{ord}_2(s) \text{ odd} \quad =: \vartheta_s \\ \varepsilon_p\left(\frac{-1}{p}\right)^{k+1}(\frac{Ms}{p},p)_2 & \text{if } s\neq 0 \text{ and } \mathrm{ord}_2(s) \text{ even,} \end{cases}$$

One can check that

$$\vartheta_s\cdot\eta_s = \varepsilon_p\left(\frac{-1}{p}\right)^k \xi_s,$$

and so

$$\mathcal{T}_{-1}(\Phi_f)(\tilde{g}_\infty) = \overline{\varepsilon_p}\sum_{s=0}^{p^2-1}\xi_s\cdot\vartheta_s\cdot\eta_s\Phi_f((A_s,1)\tilde{g}_\infty) = \left(\frac{-1}{p}\right)^k\sum_{s=0}^{p^2-1}\Phi_f((A_s,1)\tilde{g}_\infty).$$

Thus

$$q(\mathcal{T}_{-1})(f)(z) = \left(\frac{-1}{p}\right)^k\sum_{s=0}^{p^2-1} f\left(\frac{p^2z}{-4Msz+1}\right)\left(\frac{-4Msz+1}{p}\right)^{-k-1/2}$$

$$= \left(\frac{-1}{p}\right)^k\sum_{s=0}^{p^2-1} f|[(\alpha_s,\phi_{\alpha_s}(z))]_{k+1/2}(z)$$

where $\alpha_s = \begin{pmatrix} p^2 & 0 \\ -4Ms & 1 \end{pmatrix}$ and $\phi_{\alpha_s}(z) = (-4(M/p)sz+p^{-1})^{1/2}$. □

We now show that the adjoint of $q(\mathcal{T}_{-1})$ is $q(\mathcal{T}_1)$.

Consider elements $\xi = (\begin{pmatrix} 1 & 0 \\ 0 & p^2 \end{pmatrix}, p^{1/2})$ and $\eta = (\begin{pmatrix} p^2 & 0 \\ 0 & 1 \end{pmatrix}, p^{-1/2})$ in $\mathcal{G}$. For $f\in S_{k+1/2}(\Delta_0(4M))$ we have [10, Proposition 1.5]

$$f|[\Delta_0(4M)\xi\Delta_0(4M)]_{k+1/2} = U_{p^2}(f) = \left(\frac{-1}{p}\right)^k p^{(2k-3)/2}q(\mathcal{T}_1)(f).$$

We can choose β_s such that $\Gamma_0(4M)\begin{pmatrix}1 & 0\\ 0 & p^2\end{pmatrix}\Gamma_0(4M) = \bigsqcup\Gamma_0(4M)\beta_s = \bigsqcup\beta_s\Gamma_0(4M)$. So by [10, Propositions 1.1, 1.2] we have $\Delta_0(4M)\xi\Delta_0(4M) = \bigsqcup\Delta_0(4M)\xi_s = \bigsqcup\xi_s\Delta_0(4M)$ where $P(\xi_s) = \beta_s$.

Thus for $f,\ g \in S_{k+1/2}(\Gamma_0(4M))$, we have

$$\begin{aligned}&\langle f|[\Delta_0(4M)\xi\Delta_0(4M)]_{k+1/2},\ g\rangle = \langle p^{(2k-3)/2}\sum_s f|[\xi_s]_{k+1/2},\ g\rangle\\ &= \langle f,\ p^{(2k-3)/2}\sum_s g|[\xi_s^{-1}]_{k+1/2}\rangle = \langle f,\ g|[\Delta_0(4M)\eta\Delta_0(4M)]_{k+1/2}\rangle\end{aligned} \tag{1}$$

as $\Delta_0(4M)\eta\Delta_0(4M) = \bigsqcup\Delta_0(4M)\xi_s^{-1}(\begin{pmatrix}p^2 & 0\\ 0 & p^2\end{pmatrix}, 1)$ and elements of type $(aI, 1)$ belongs to the center of $\mathcal{G}$ and acts trivially via slash operator.

Using triangular decomposition we check that $\Gamma_0(4M)\begin{pmatrix}p^2 & 0\\ 0 & 1\end{pmatrix}\Gamma_0(4M) = \bigsqcup_{s=0}^{p^2-1}\Gamma_0(4M)\begin{pmatrix}p^2 & 0\\ 0 & 1\end{pmatrix}\begin{pmatrix}1 & 0\\ -4Ms & 1\end{pmatrix}$ and so

$$\Delta_0(4M)\eta\Delta_0(4M) = \bigsqcup_{s=0}^{p^2-1}\Delta_0(4M)\ \eta\ (\begin{pmatrix}1 & 0\\ -4Ms & 1\end{pmatrix}, (-4Msz+1)^{1/2})$$

$$= \bigsqcup_{s=0}^{p^2-1}\Delta_0(4M)(\begin{pmatrix}p^2 & 0\\ -4Ms & 1\end{pmatrix}, (-4(M/p)sz+p^{-1})^{1/2}).$$

Hence $g|[\Delta_0(4M)\eta\Delta_0(4M)]_{k+1/2} = \left(\frac{-1}{p}\right)^k p^{(2k-3)/2}q(\mathcal{T}_{-1})(g)$. Thus by equation (1) above we obtain the following

Proposition 4.6. *The operator $q(\mathcal{T}_{-1})$ is adjoint of $q(\mathcal{T}_1)$ and consequently $\widetilde{Q}_p$ is self-adjoint with respect to Petersson inner product.*

5. Local Hecke algebra at prime 2 and Kohnen's plus space

Loke and Savin [6] studied genuine local Hecke algebra of $\widetilde{\mathrm{SL}}(\mathbb{Q}_2)$ with respect to $\overline{K_0^2(4)}$ and central character of M_2 and gave an interpretation of Kohnen's plus space at level 4 in terms of certain elements in this 2-adic Hecke algebra. In this section we shall recall their work and describe Kohnen's plus space at level $4M$ for M odd in a similar way.

Let γ be a genuine character of $\overline{K_0^2(4)}$ determined by its value on $(-I, 1)$ such that it is trivial on $K_1^2(4)$. This character can be extended to the normalizer $N_{\widetilde{\mathrm{SL}}(\mathbb{Q}_2)}(T)$ by defining $\gamma((h(2^n), 1)) = 1$ for all integers n and $\gamma((w(1), 1)) = (1+\gamma((-I, 1)))/\sqrt{2} =: \zeta_8$, a primitive 8th root of unity. Let $H(\overline{K_0^2(4)}, \gamma)$ be genuine Hecke algebra of $\widetilde{\mathrm{SL}}(\mathbb{Q}_2)$ with respect to $\overline{K_0^2(4)}$ and γ. Let $\mathcal{T}_n = X_{(h(2^n),1)}$, $\mathcal{U}_n = X_{(w(2^{-n}),1)} \in H(\overline{K_0^2(4)}, \gamma)$ (defined as in the odd prime case).

Theorem 2. *(Loke-Savin [6]) For $m, n \in \mathbb{Z}$,*

(1) If $mn \geq 0$ then $\mathcal{T}_m * \mathcal{T}_n = \mathcal{T}_{m+n}$.
(2) $\mathcal{U}_1 * \mathcal{T}_n = \mathcal{U}_{n+1}$ and $\mathcal{T}_n * \mathcal{U}_1 = \mathcal{U}_{1-n}$.
(3) $\mathcal{U}_1 * \mathcal{U}_n = \mathcal{T}_{n-1}$ and $\mathcal{U}_n * \mathcal{U}_1 = \mathcal{T}_{1-n}$.

The Hecke algebra $H(\overline{K_0^2(4)}, \gamma)$ is generated by $\mathcal{U}_0$ and $\mathcal{U}_1$ modulo relations $(\mathcal{U}_0 - 2\sqrt{2})(\mathcal{U}_0 + \sqrt{2}) = 0$ and $\mathcal{U}_1^2 = 1$.

In particular, note from the above that $\mathcal{U}_2 = \mathcal{U}_1 * \mathcal{T}_1 = \mathcal{U}_1 * \mathcal{U}_0 * \mathcal{U}_1$ and so $(\mathcal{U}_2 - 2\sqrt{2})(\mathcal{U}_2 + \sqrt{2}) = 0$.

Let χ be the trivial character modulo 4, thus $\chi_0 = \left(\frac{-1}{\cdot}\right)^k$. Let $\gamma((-I, 1)) = -i^{2k+1}$. Then, for any $k_0 = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K_0^2(4)$ we can check that $\tilde{\epsilon}_2(k_0)\chi_{0,2}(d) = \gamma((k_0, 1))$.

Proposition 5.1. *(Loke-Savin [6]) For $f \in S_{k+1/2}(\Gamma_0(4), \chi)$,*

(1) $q(\mathcal{T}_1)(f)(z) = 2^{(3-2k)/2} U_4(f)(z)$.
(2) $q(\mathcal{U}_1)(f)(z) = \left(\frac{2}{2k+1}\right) W_4(f)(z)$ where the operator W_4 is given by $W_4(f)(z) = (-2iz)^{-k-1/2} f(-1/4z)$.

Niwa [7] considered operator $R = W_4 U_4$ on $S_{k+1/2}(\Gamma_0(4), \chi)$, proved that it is self-adjoint and that $(R - \alpha_1)(R - \alpha_2) = 0$ where $\alpha_1 = \left(\frac{2}{2k+1}\right) 2^k$, $\alpha_2 = -\frac{\alpha_1}{2}$. Kohnen [4] defined his plus space $S^+(4)$ at level 4 to be the α_1-eigenspace of R in $S_{k+1/2}(\Gamma_0(4))$. It follows from the above proposition that $S^+(4)$ is the 2-eigenspace of $q(\mathcal{U}_1)q(\mathcal{T}_1)/\sqrt{2}$ and hence that of $q(\mathcal{U}_2)/\sqrt{2}$.

In the case level is $4M$ with M odd and χ is a trivial character modulo $4M$, Kohnen [5] defines a classical operator Q on $S_{k+1/2}(\Gamma_0(4M))$ in order to obtain his plus space. The operator Q is defined by

$$Q := [\Delta_0(4M, \chi)\rho\Delta_0(4M, \chi)] \text{ where } \rho = (\begin{pmatrix} 4 & 1 \\ 0 & 4 \end{pmatrix}, e^{\pi i/4}).$$

By [5, Proposition 1] Q is self-adjoint and satisfies $(Q-\alpha)(Q-\beta) = 0$ where $\alpha = (-1)^{[(k+1)/2]} 2\sqrt{2}$, $\beta = -\alpha/2$, and the plus space $S^+_{k+1/2}(4M)$ is precisely the α-eigenspace of Q.

In our work we need that Kohnen's plus space at level $4M$ is the 2-eigenspace of the product $q(\mathcal{U}_1)q(\mathcal{T}_1)/\sqrt{2}$. We prove this below.

Proposition 5.2. *Let $f \in S_{k+1/2}(\Gamma_0(4M))$ with M odd. Then we have*

$$Q(f) = \left(\frac{2}{2k+1}\right) q(\mathcal{U}_2)(f) = \left(\frac{2}{2k+1}\right) q(\mathcal{U}_1)q(\mathcal{T}_1)(f).$$

Consequently $S^+_{k+1/2}(4M)$ is the 2-eigenspace of $q(\mathcal{U}_1)q(\mathcal{T}_1)/\sqrt{2}$.

Proof. Following [5, Proposition 1] we can write

$$Q(f) = \sum_{s=0}^{4} f|[\rho]_{k+1/2}|[\begin{pmatrix} 1 & 0 \\ 4Ms & 1 \end{pmatrix}, (4Msz+1)^{1/2}]_{k+1/2}$$
$$= e^{-(2k+1)\pi i/4} \sum_{s=0}^{4} f|[\begin{pmatrix} 4+4Ms & 1 \\ 16Ms & 4 \end{pmatrix}, (4Msz+1)^{1/2}]_{k+1/2}.$$

and it's adjoint

$$\tilde{Q}(f) = \sum_{s=0}^{4} f|[\begin{pmatrix} 4 & -1 \\ 0 & 4 \end{pmatrix}, e^{-\pi i/4}]_{k+1/2}|[\begin{pmatrix} 1 & 0 \\ 4Ms & 1 \end{pmatrix}, (4Msz+1)^{1/2}]_{k+1/2}$$
$$= e^{(2k+1)\pi i/4} \sum_{s=0}^{4} f|[\begin{pmatrix} 4-4Ms & -1 \\ 16Ms & 4 \end{pmatrix}, (4Msz+1)^{1/2}]_{k+1/2}.$$

Since Q is self adjoint, $Q = \tilde{Q}$.

We now compute $q(\mathcal{U}_2)(f)(z)$ for $f \in S_{k+1/2}(4M)$.

Let $\tilde{g}_\infty \in \widetilde{\mathrm{SL}_2}(\mathbb{R})$ be such that $\tilde{g}_\infty i = z$. Since $K_0^2(4)w(2^{-2})K_0^2(4) = \bigcup_{s\in\mathbb{Z}/4\mathbb{Z}} y(4M(1-s))w(2^{-2})K_0^2(4)$,

$$\mathcal{U}_2(\Phi_f)(\tilde{g}_\infty) = \overline{\zeta_8} \sum_{s=0}^{3} \Phi_f(\tilde{g}_\infty(y(4M(1-s)),1)(w(2^{-2}),1)).$$

Take $A_s = \begin{pmatrix} 1-\left(\frac{-1}{M}\right)Ms & -\left(\frac{-1}{M}\right)/4 \\ 4Ms & 1 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Q})$, so $s_{\mathbb{Q}}(A_s) = (A_s, 1)$. The ∞-component of

$$s_{\mathbb{Q}}(A_s)\ \tilde{g}_\infty\ (y(4M(1-s)),1)(w(2^{-2}),1))$$

is $(A_s,1)\tilde{g}_\infty$, for a prime q such that $(q,2M)=1$ the q-component is $(A_s,1) \in \mathrm{SL}_2(\mathbb{Z}_q)\times\{1\}$, for an odd prime p such that $p^b \| M$, the p-component is $(A_s,1) \in K_0^p(p^b)\times\{1\}$ while the 2-component is

$$(A_s,1)(y(4M(1-s)),1)(w(2^{-2}),1)) = (\begin{pmatrix} \left(\frac{-1}{M}\right) & \frac{1-M\left(\frac{-1}{M}\right)}{4} \\ -4 & M \end{pmatrix}, 1).$$

Since M is odd, it is clear that $\frac{1-M\left(\frac{-1}{M}\right)}{4} \in \mathbb{Z}_2$ and so the 2-component is in $K_0^2(4)\times\{1\}$. The p-component acts trivially while the 2-component acts by $(\tilde{\gamma_2}(M))^{-1}(-1,M)_2\chi_{0,2}(M) =: \omega_M$. Hence

$$q(\mathcal{U}_2)(f)(z) = \overline{\zeta_8}\ \omega_M \sum_{s=0}^{3} f(A_s z)J(A_s,z)^{-2k-1}$$
$$= \overline{\zeta_8}\ \omega_M \sum_{s=0}^{3} f\left(\frac{(4-4M\left(\frac{-1}{M}\right)sz) - \left(\frac{-1}{M}\right)}{16Msz+4}\right)(4Msz+1)^{-k-1/2}$$

We note that $e^{(2k+1)\pi i/4} = \left(\frac{2}{2k+1}\right)\overline{\zeta_8}$. Thus when $M \equiv 1 \pmod 4$ since $\omega_M = 1$, comparing the expression of $\tilde{Q}$ and $q(\mathcal{U}_2)$ we see that $\tilde{Q}(f) = \left(\frac{2}{2k+1}\right) q(\mathcal{U}_2)(f)$. In the case $M \equiv 3 \pmod 4$ we get that $\omega_M = -i(-1)^k$, so $\left(\frac{2}{2k+1}\right)\overline{\zeta_8}\omega_M = e^{-(2k+1)\pi i/4}$ and consequently $Q(f) = \left(\frac{2}{2k+1}\right) q(\mathcal{U}_2)(f)$. Hence we are done.

The last statement follows since $(-1)^{[(k+1)/2]} = \left(\frac{2}{2k+1}\right)$. □

Following above calculations we can translate $\mathcal{T}_1,\ \mathcal{U}_1,\ \mathcal{U}_0 \in H(\overline{K_0^2(4)}, \gamma)$ to classical operators on $S_{k+1/2}(\Gamma_0(4M))$.

Proposition 5.3. *For $f \in S_{k+1/2}(\Gamma_0(4M))$,*

(1) $q(\mathcal{T}_1)(f)(z) = 2^{(3-2k)/2} U_4(f)(z)$.

(2) $q(\mathcal{U}_1)(f)(z) = \overline{\zeta_8}\left(\frac{2}{M}\right)\left(\frac{-1}{M}\right)^{k+3/2} f|[W, \phi_W(z)]_{k+1/2}(z)$ *where* $W = \begin{pmatrix} 4n & m \\ 4M & 2 \end{pmatrix}$ *with $a, b \in \mathbb{Z}$ such that $2n - mM = 1$ and $\phi_W(z) = (2Mz+1)^{1/2}$.*

(3) $q(\mathcal{U}_0)(f)(z) = \overline{\zeta_8}\left(\frac{-1}{M}\right)^{k+3/2} \sum_{s=0}^{3} f|[A_s, \phi_{A_s}(z)]_{k+1/2}(z)$ *where* $A_s = \begin{pmatrix} n & -ns+m \\ M & -Ms+4 \end{pmatrix}$ *with $m, n \in \mathbb{Z}$ such that $4n - mM = 1$ and $\phi_W(z) = (Mz + 4 - Ms)^{1/2}$.*

Define $\widetilde{Q}_2 := q(\mathcal{U}_0)/\sqrt{2} = q(\mathcal{T}_1)q(\mathcal{U}_1)/\sqrt{2}$ and $\widetilde{W}_4 := q(\mathcal{U}_1)$. Further define $\widetilde{Q}_2'$ to be the conjugate of $\widetilde{Q}_2$ by $\widetilde{W}_4$, so $\widetilde{Q}_2' = q(\mathcal{U}_1)q(\mathcal{T}_1)/\sqrt{2}$. Then the Kohnen's plus space at level $4M$ is the 2-eigenspace of $\widetilde{Q}_2'$.

6. The minus space of half-integral weight forms

Let M be odd and square-free. In this section we shall define the minus space $S^-_{k+1/2}(4M)$ of weight $k+1/2$ and level $4M$ and show that there is an Hecke algebra isomorphism between$S^-_{k+1/2}(4M)$ and $S^{\text{new}}_{2k}(\Gamma_0(2M))$.

We shall first start with defining the minus space at level 4 and then at level $4p$ for p an odd prime. We can then extend our definition to level $4M$ where M is as above.

6.1. Eigenvalues of U_p.

Let N be a positive integer and p be a prime strictly dividing N, i.e. $p \| N$. We compute the eigenvalues of the Hecke operator U_p on the space of old forms of weight $2k$ on $\Gamma_0(N)$.

For every positive integer n and a modular form F we denote by

$$F_n(z) := V(n)F(z) = F(nz).$$

Recall that $V(n)$ is the shift operator sending $S_{2k}(\Gamma_0(L))$ to $S_{2k}(\Gamma_0(nL))$.

Let M be a positive integer such that $p \nmid M$. Let $F \in S_{2k}(\Gamma_0(M))$. Then by well-known action of T_p and U_p we have

$$U_p(F)(z) = T_p(F)(z) - p^{2k-1}F_p(z). \tag{2}$$

Assume that $F \in S_{2k}(\Gamma_0(M))$ is a primitive Hecke eigenform and a_p is the p-th Fourier coefficient of F. Then $T_p(F) = a_pF$. It is known that a_p is real and by the Ramanujan conjecture proved by Deligne we have that $|a_p| \leq 2 \cdot p^{(2k-1)/2}$.

Lemma 6.1. *(a) If $(p, n) = 1$ then $U_p(F_n) = a_pF_n - p^{2k+1}F_{np}$.*
(b) If $p \mid n$ then $U_p(F_n) = F_{n/p}$.

Proof. (a) It is well known that if $(p, n) = 1$ then $V(n)T_p(F) = T_pV(n)F$. Hence using (2) and that F is a primitive Hecke eigenform we get that

$$\begin{aligned} U_p(F_n) &= T_p(F_n) - p^{2k-1}F_{np} = V(n)T_p(F) - p^{2k-1}F_{np} \\ &= V(n)a_pF - p^{2k-1}F_{np} = a_pF_n - p^{2k-1}F_{np}. \end{aligned}$$

For (b) write $n = mp$. Then

$$U_p(F_n)(z) = \frac{1}{p}\sum_{k=0}^{p-1} F_{mp}\left(\frac{z+k}{p}\right) = \frac{1}{p}\sum_{k=0}^{p-1} F_m(z+k) = F_{n/p}(z)$$

□

Thus U_p stabilizes the two dimensional subspace spanned by F_n and F_{np} for $(p, n) = 1$. We will compute the eigenvalues of U_p on this space. If $M = \lambda F_n + \beta F_{np}$ is an eigenfunction of U_p then it follows from (2) that $\lambda \neq 0$. Hence we can assume that $\lambda = 1$. We have

$$U_p(F_n + \beta F_{np}) = (a_p + \beta)F_n - p^{2k-1}F_{np}$$

It is clear from above that β cannot be zero and that M is an eigenfunction if and only if $a_p + \beta = -p^{2k-1}/\beta$ with eigenvalue $a_p + \beta$. Hence $\beta^2 + a_p\beta + p^{2k-1} = 0$ and we have

$$\beta = \frac{-a_p \pm \sqrt{a_p^2 - 4p^{2k-1}}}{2}.$$

The eigenvalues of U_p on the subspace $\langle F_n, F_{np}\rangle$ are

$$a_p + \beta = \frac{a_p \pm \sqrt{a_p^2 - 4p^{2k-1}}}{2}.$$

Proposition 6.2. *The eigenvalues of $(U_p)^2$ on the two dimensional subspace spanned by F_n and F_{np} are either complex or $\pm p^{2k-1}$.*

Proof. Using the Ramanujan conjecture we can see that the eigenvalues of U_p are real or purely imaginary if and only if $a_p = \pm 2p^{k-1/2}$ or $a_p = 0$. In those cases the eigenvalue of $(U_p)^2$ are precisely $\pm p^{2k-1}$. □

6.2. **Minus space for $\Gamma_0(4)$.** We recall the following theorem of Niwa which was obtained by proving equality of traces of Hecke operators.

Theorem 3. *(Niwa [7]) Let M be odd and square-free. There exists an isomorphism of vector spaces $\psi : S_{k+1/2}(\Gamma_0(4M)) \to S_{2k}(\Gamma_0(2M))$ satisfying*

$$T_p(\psi(f)) = \psi(T_{p^2}(f)) \quad \text{for all primes } p \text{ coprime to } 2M.$$

Moreover if $f \in S_{k+1/2}(\Gamma_0(4))$ then we further have

$$U_2(\psi(f)) = \psi(U_4(f)), \quad T_p(\psi(f)) = \psi(T_{p^2}(f)) \text{ for all odd primes } p.$$

We also recall Shimura lift [10]: For t a positive square-free integer, there is a linear map $\mathrm{Sh}_t : S_{k+1/2}(\Gamma_0(4M)) \to S_{2k}(\Gamma_0(2M))$ given by

$$\mathrm{Sh}_t\left(\sum_{n=1}^{\infty} a_n q^n\right) = \sum_{n=1}^{\infty}\left(\sum_{d|n}\left(\frac{-1}{d}\right)^k\left(\frac{t}{d}\right)d^{k-1}a\left(t\frac{n^2}{d^2}\right)\right)q^n.$$

We note the following observations [8]:
(a) Sh_t need not be $1-1$ but $\mathrm{Sh}_t(f) = 0$ for all square-free t implies $f = 0$.
(b) Sh_t commutes with all Hecke operators, i.e., $T_p(\mathrm{Sh}_t(f)) = \mathrm{Sh}_t(T_{p^2}(f))$ for all primes p coprime to $2M$ and $U_p(\mathrm{Sh}_t(f)) = \mathrm{Sh}_t(U_{p^2}(f))$ for all primes p dividing $2M$.

We need the following theorem of Kohnen.

Theorem 4. *(Kohnen [4])*

1. $\dim(S^+(4)) = \dim(S_{2k}(\Gamma_0(1)))$.
2. *$S^+(4)$ has a basis of eigenforms for all the operators T_{p^2}, p odd.*
3. *If f is such an eigenform then $\psi(f)$ is an old form and $\psi(f) = \lambda F + \beta F_2$ where $F \in S_{2k}(\Gamma_0(1))$ is a primitive eigenform determined by the eigenvalues of f.*

Define $A^+(4) = \widetilde{W}_4 S^+(4)$. We know that $S^+(4)$ is the 2-eigenspace of $\widetilde{Q}_2'$, hence $A^+(4)$ is the 2-eigenspace of $\widetilde{Q}_2$. Following above theorem of Kohnen we have $\dim(A^+(4)) = \dim(S_{2k}(\Gamma_0(1))$ and

Corollary 6.3.
1. *$A^+(4)$ has a basis of eigenforms under T_{p^2} for all p odd.*
2. *ψ maps $A^+(4)$ into the space of old forms in $S_{2k}(\Gamma_0(2))$.*

Proof. Let $f \in S^+(4)$ be an eigenform under T_{p^2} for all p odd satisfying $T_{p^2}(f) = \lambda_p f$. Since $\widetilde{W}_4$ commutes with all such T_{p^2}, we get that $g = \widetilde{W}_4 f \in A^+(4)$ is also an eigenform under all T_{p^2} with eigenvalues λ_p. By Theorem 3, $\psi(f)$ and $\psi(g)$ are eigenforms in $S_{2k}(\Gamma_0(2))$ under all T_p with the same set of eigenvalues λ_p. Since $\psi(f)$ is an old form it follows from Atkin-Lehner [1] that $\psi(g)$ is also an old form (belonging to the same two dimensional subspace $\langle F, F_2\rangle$). □

Proposition 6.4. $S^+(4) \bigcap A^+(4) = \{0\}$.

Proof. Suppose there is a nonzero $f \in S^+(4) \bigcap A^+(4)$. We can assume that f is an eigenform under T_{p^2} for all p odd (as such T_{p^2} stabilizes the intersection $S^+(4) \bigcap A^+(4)$). Then $\widetilde{Q}_2(f) = 2f = \widetilde{Q}'_2(f)$. This implies that $U_4\widetilde{W}_4(f) = 2^k f = \widetilde{W}_4 U_4(f)$. Thus

$$(U_4)^2(f) = 2^k U_4 \widetilde{W}_4(f) = 2^{2k} f.$$

Applying ψ to the above equation we get that $(U_2)^2(\psi(f)) = 2^{2k}\psi(f)$. Now $\psi(f) \in \langle F, F_2 \rangle$ for some primitive form $F \in S_{2k}(\Gamma_0(1))$ and by Proposition 6.2, the eigenvalues of $(U_2)^2$ on this subspace are either complex or 2^{2k-1}. This is a contradiction. □

Define $S^-_{k+1/2}(4)$ to be the orthogonal complement of $S^+(4) \oplus A^+(4)$. That is, $S^-_{k+1/2}(4)$ is the common eigenspace with the eigenvalue -1 of the operators $\widetilde{Q}_2$ and $\widetilde{Q}'_2$. We shall write $S^-_{k+1/2}(4)$ simply by $S^-(4)$. So we have

$$S_{k+1/2}(\Gamma_0(4)) = S^+(4) \oplus A^+(4) \oplus S^-(4) \tag{3}$$

Theorem 5. *$S^-(4)$ has a basis of eigenforms for all the operators T_{p^2}, p odd; these eigenforms are also eigenfunctions under U_4. If two eigenforms in $S^-(4)$ share the same eigenvalues for all T_{p^2} then they are a scalar multiple of each other. ψ induces a Hecke algebra isomorphism:*

$$S^-_{k+1/2}(4) \cong S^{\text{new}}_{2k}(\Gamma_0(2)).$$

Proof. Since ψ maps $S^+(4) \oplus A^+(4)$ into $S^{\text{old}}_{2k}(\Gamma_0(2))$ and $\dim(S^+(4) \oplus A^+(4))$ $= 2\dim(S_{2k}(\Gamma_0(1))) = \dim(S^{\text{old}}_{2k}(\Gamma_0(2)))$, we get that ψ maps this direct sum onto $S^{\text{old}}_{2k}(\Gamma_0(2))$.

Now T_{p^2} commutes with $\widetilde{Q}_2$ and $\widetilde{Q}'_2$ for every odd prime p so we get that T_{p^2} stabilizes $S^-(4)$, hence it has a basis of eigenforms for all T_{p^2} with p odd.

If f is such an eigenform then $F := \psi(f)$ is an eigenform in $S_{2k}(\Gamma_0(2))$ under all T_p, p odd. By Atkin-Lehner [1] F is either an old form or a newform. Since ψ is injective, it follows that F must be a newform. So ψ maps the space $S^-(4)$ into the space $S^{\text{new}}_{2k}(\Gamma_0(2))$. By equality of dimensions, we get that ψ is an isomorphism of $S^-(4)$ onto $S^{\text{new}}_{2k}(\Gamma_0(2))$. Consequently by [1] an eigenform in $S^-(4)$ under all T_{p^2} for p odd is uniquely determined up to scalar multiplication.

Further for such an eigenform f, by [1, Theorem 3], $U_2(F) = -2^{k-1}\lambda(2)F$ where $\lambda(2) = \pm 1$. Thus $\psi(U_4(f)) = U_2(F) \in S^{\text{new}}_{2k}(\Gamma_0(2))$, so $U_4(f)$ belongs to $S^-(4)$. Since U_4 commutes with T_{p^2} for all p odd, we get that $U_4(f)$ is an eigenform under all T_{p^2} with the same eigenvalues as f and hence is a scalar multiple of f. □

6.3. **Minus space for $\Gamma_0(4p)$ for p an odd prime.** In this subsection we need the involution $\widetilde{W}_{p^2}$, the operators U_{p^2}, $\widetilde{Q}_p$ and $\widetilde{Q}'_p = \widetilde{W}_{p^2}\widetilde{Q}_p\widetilde{W}_{p^2}$ on $S_{k+1/2}(\Gamma_0(4p))$ that we defined in Section 4.

Consider the subspace $\mathcal{V}(1)$ of $S_{2k}(\Gamma_0(2p))$ coming from the old forms at level 1, that is,

$$\mathcal{V}(1) = S_{2k}(\Gamma_0(1)) \oplus V(2)S_{2k}(\Gamma_0(1)) \oplus V(p)S_{2k}(\Gamma_0(1)) \oplus V(2p)S_{2k}(\Gamma_0(1)).$$

We consider the eigenvalues of $(U_p)^2$ on $\mathcal{V}(1)$.

Lemma 6.5. *The operator U_p stabilizes $\mathcal{V}(1)$. If an eigenvalue λ of $(U_p)^2$ on this space is real then $\lambda = \pm p^{2k-1}$.*

Proof. For a a primitive Hecke eigenform F in $S_{2k}(\Gamma_0(1))$ consider the four dimensional subspace spanned by F, F_2, F_p, F_{2p}. Then $\mathcal{V}(1)$ is a direct sum of such four dimensional subspaces. By Lemma 6.1, U_p preserves the two dimensional subspaces spanned by F and F_p and the two dimensional subspace spanned by F_2 and F_{2p}. It follows by Proposition 6.2, that the eigenvalues of $(U_p)^2$ on these two dimensional subspaces are either complex or $\pm p^{2k-1}$. □

Let $R := S^+(4) \oplus A^+(4)$. Then we have

Proposition 6.6. $R \bigcap \widetilde{W}_{p^2} R = \{0\}$

Proof. Let $f \neq 0$ belongs to the intersection. We can again assume that f is an eigenform under T_{q^2} for all primes q coprime to $2p$. Since by Corollary 4.3(4), $S_{k+1/2}(\Gamma_0(4))$ is contained in the p eigenspace of $\widetilde{Q}_p$ and so $\widetilde{W}_{p^2} S_{k+1/2}(\Gamma_0(4))$ is contained in the p-eigenspace of $\widetilde{Q}'_p$ we have $\widetilde{Q}_p(f) = pf = \widetilde{Q}'_p(f)$. Using $\widetilde{Q}_p = \left(\frac{-1}{p}\right)^k p^{1-k} U_{p^2} \widetilde{W}_{p^2}$ in the above equality we obtain

$$(U_{p^2})^2(f) = p^{2k} f.$$

Since $f \neq 0$, there exists a t square-free such that the Shimura lift $\mathrm{Sh}_t(f) \neq 0$. Applying this Sh_t to the above equation we get that $(U_p)^2(\mathrm{Sh}_t(f)) = p^{2k}\mathrm{Sh}_t(f)$. Since Sh_t commutes with all the Hecke operators we get that $\mathrm{Sh}_t(f) \in \mathcal{V}(1)$. But by Lemma 6.5, the eigenvalues of $(U_p)^2$ on $\mathcal{V}(1)$ are either complex or p^{2k-1} leading to a contradiction. □

Corollary 6.7. *Niwa's map ψ maps $R \oplus \widetilde{W}_{p^2} R$ isomorphically onto $\mathcal{V}(1)$.*

Proof. As before (see Corollary 6.3(2)) ψ maps $R \oplus \widetilde{W}_{p^2} R$ into $\mathcal{V}(1)$. It follows from the equality of dimensions that the map is onto. □

Next we consider the following subspace of $S_{2k}(\Gamma_0(2p))$ coming from the old forms at level 2,

$$\mathcal{V}(2) = S_{2k}^{\mathrm{new}}(\Gamma_0(2)) \oplus V(p)S_{2k}^{\mathrm{new}}(\Gamma_0(2)).$$

This space is a direct sum of two dimensional subspaces spanned by F and F_p where F is a primitive Hecke eigenform in $S_{2k}^{\mathrm{new}}(\Gamma_0(2))$. Using Proposition 6.2 we have the following lemma.

Lemma 6.8. *If an eigenvalue λ of $(U_p)^2$ on $\mathcal{V}(2)$ is real then $\lambda = \pm p^{2k-1}$.*

Since (by Theorem 5) ψ maps $S^-_{k+1/2}(4)$ isomorphically onto $S^{\text{new}}_{2k}(\Gamma_0(2))$, it follows that ψ maps $\widetilde{W}_{p^2}S^-_{k+1/2}(4)$ into the space $\mathcal{V}(2)$. The proof of the following is identical to that of Proposition 6.6.

Proposition 6.9. $S^-_{k+1/2}(4)\bigcap\widetilde{W}_{p^2}S^-_{k+1/2}(4)=\{0\}$.

Corollary 6.10. *ψ maps $S^-_{k+1/2}(4)\oplus\widetilde{W}_{p^2}S^-_{k+1/2}(4)$ isomorphically onto $\mathcal{V}(2)$.*

Finally, we consider the following subspace of $S_{2k}(\Gamma_0(2p))$ coming from the old forms at level p,

$$\mathcal{V}(p)=S^{\text{new}}_{2k}(\Gamma_0(p))\oplus V(2)S^{\text{new}}_{2k}(\Gamma_0(p)).$$

This space is a direct sum of two dimensional subspaces spanned by F and F_2 where F is a primitive Hecke eigenform in $S^{\text{new}}_{2k}(\Gamma_0(p))$. We have

Lemma 6.11. *If an eigenvalue λ of $(U_2)^2$ on $\mathcal{V}(p)$ is real then $\lambda=\pm 2^{2k-1}$.*

Let $S^{+,\text{new}}_{k+1/2}(4p)$ be the new space inside the plus space in $S_{k+1/2}(\Gamma_0(4p))$. Kohnen [5, Theorem 2] proved that ψ maps $S^{+,\text{new}}_{k+1/2}(4p)$ into $\mathcal{V}(p)$ and the dimension of $S^{+,\text{new}}_{k+1/2}(4p)$ equals the dimension of $S^{\text{new}}_{2k}(\Gamma_0(p))$. Then as before ψ maps $\widetilde{W}_4S^{+,\text{new}}_{k+1/2}(4)$ into the space $\mathcal{V}(p)$ and we have the following proposition and corollary.

Proposition 6.12. $S^{+,\text{new}}_{k+1/2}(4p)\bigcap\widetilde{W}_4S^{+,\text{new}}_{k+1/2}(4p)=\{0\}$.

Corollary 6.13. *ψ maps $S^{+,\text{new}}_{k+1/2}(4p)\oplus\widetilde{W}_4S^{+,\text{new}}_{k+1/2}(4p)$ isomorphically onto $\mathcal{V}(p)$.*

We define the following subspace of $S_{k+1/2}(\Gamma_0(4p))$,

$$E=R\oplus\widetilde{W}_{p^2}R\oplus S^-_{k+1/2}(4)\oplus\widetilde{W}_{p^2}S^-_{k+1/2}(4)\oplus S^{+,\text{new}}_{k+1/2}(4p)\oplus\widetilde{W}_4S^{+,\text{new}}_{k+1/2}(4p).$$

By Corollary 6.7, 6.10 and 6.13, we get that ψ maps the space E isomorphically onto the old space $S^{\text{old}}_{2k}(\Gamma_0(2p))$. We define the minus space to be the orthogonal complement of E under the Petersson inner product. That is,

$$S^-_{k+1/2}(4p):=E^{\perp}.$$

Theorem 6. *$S^-_{k+1/2}(4p)$ has a basis of eigenforms for all the operators T_{q^2}, q an odd prime different than p, uniquely determined up to a non-zero scalar multiplication. ψ maps the space $S^-_{k+1/2}(4p)$ isomorphically to the space $S^{\text{new}}_{2k}(\Gamma_0(2p))$.*

Proof. Since the operators T_{q^2} with $(q,2p)=1$ stabilize the space E and since they are self adjoint with respect to the Petersson inner product, it follows that they stabilize the space $S^-_{k+1/2}(4p)$, hence it has a basis of eigenforms for all such operators T_{q^2}. If f is such an eigenform then $\psi(f)\in S_{2k}(\Gamma_0(2p))$

is also an eigenform for all the operators T_q, $(q, 2p) = 1$ and thus (by [1]) $\psi(f)$ is either an old form or a newform. Since ψ is injective and maps E onto $S_{2k}^{\text{old}}(\Gamma_0(2p))$, it follows that $\psi(f)$ is a newform. Thus ψ maps the space $S_{k+1/2}^{-}(4p)$ into the space $S_{2k}^{\text{new}}(\Gamma_0(2p))$. By equality of dimensions, we get that ψ maps the space $S_{k+1/2}^{-}(4p)$ isomorphically onto $S_{2k}^{\text{new}}(\Gamma_0(2p))$. Consequently an eigenform in $S_{k+1/2}^{-}(4p)$ is uniquely determined up to a scalar multiplication. □

Corollary 6.14. *Let $f \in S_{k+1/2}^{-}(4p)$ be a Hecke eigenform for all the operators T_{q^2}, q prime and $(q, 2p) = 1$. Then $\widetilde{W}_{p^2} f = \beta(p) f$, $\widetilde{W}_4 f = \beta(2) f$ where $\beta(p) = \pm 1$, $\beta(2) = \pm 1$.*

Proof. Let $g = \widetilde{W}_{p^2} f$. Since $\widetilde{W}_{p^2}$ commutes with all the operators T_{q^2} for $(q, 2p) = 1$ we get that g is an eigenform for all the operators T_{q^2} with the same eigenvalues as f. Since $\psi(f)$ is a newform, it follows [1] that $\psi(g)$ is a scalar multiple of $\psi(f)$. Since ψ is an isomorphism we get that g is a scalar multiple of f. Since $\widetilde{W}_{p^2}$ is an involution we get that the scalar is ± 1. The same proof applies to $\widetilde{W}_4$. □

Let $f \in S_{k+1/2}^{-}(4p)$ be a Hecke eigenform for all the operators T_{q^2} as above. It follows that $F := \psi(f)$ is a Hecke eigenform in $S_{2k}^{\text{new}}(\Gamma_0(2p))$ for all the operators T_q, $(q, 2p) = 1$. Since the Shimura lift $\text{Sh}_t(f)$ is also an eigenform for all the operators T_q with the same eigenvalues as F, it follows from [1] that $\text{Sh}_t(f)$ is a scalar multiple of F (which could be zero). Also, $U_p(F) = -p^{k-1}\lambda(p)F$ where $\lambda(p) = \pm 1$ and $U_2(F) = -2^{k-1}\lambda(2)F$ where $\lambda(2) = \pm 1$.

Proposition 6.15. *Let $f \in S_{k+1/2}^{-}(4p)$ be a Hecke eigenform for all the operators T_{q^2}, q prime and $(q, 2p) = 1$. Then*

$$U_{p^2}(f) = -p^{k-1}\lambda(p) f, \quad U_4(f) = -2^{k-1}\lambda(2) f$$

where $\lambda(p) = \pm 1$ and $\lambda(2) = \pm 1$ are defined as above.

Proof. Let $g = U_{p^2} f$. Then $\text{Sh}_t(g) = U_p \text{Sh}_t(f) = -p^{k-1}\lambda(p)\text{Sh}_t(f)$ for every positive square-free integer t. It follows that $\text{Sh}_t(g - p^{k-1}\lambda(p) f) = 0$ for all such t implying $g - p^{k-1}\lambda(p) f = 0$ which is what we need. For the prime 2, the proof is the same. □

Proposition 6.16. *Let $f \in S_{k+1/2}^{-}(4p)$. Then $\widetilde{Q}_p(f) = -f = \widetilde{Q}'_p(f)$ and $\widetilde{Q}_2(f) = -f = \widetilde{Q}'_2(f)$.*

Proof. Let $f \in S_{k+1/2}^{-}(4p)$ be a Hecke eigenform for all the operators T_{q^2}, $(q, 2p) = 1$. Since $\widetilde{Q}_p = \left(\frac{-1}{p}\right)^k p^{1-k} U_{p^2} \widetilde{W}_{p^2}$ and $\widetilde{Q}_2 = 2^{1-k} U_4 \widetilde{W}_4$ it follows from Corollary 6.14 and Proposition 6.15 that f is an eigenform for the operators $\widetilde{Q}_p$, $\widetilde{Q}'_p$, $\widetilde{Q}_2$ and $\widetilde{Q}'_2$ with eigenvalues ± 1. However, the eigenvalues

of $\widetilde{Q}_p$, $\widetilde{Q}'_p$ are p and -1 and the eigenvalues of $\widetilde{Q}_2$ and $\widetilde{Q}'_2$ are 2 and -1 hence the eigenvalues have to be -1. Since $S^-_{k+1/2}(4p)$ has a basis of such eigenforms we get the result. □

Theorem 7. *Let $f \in S_{k+1/2}(4p)$. Then $f \in S^-_{k+1/2}(4p)$ if and only if $\widetilde{Q}_p(f) = -f = \widetilde{Q}'_p(f)$ and $\widetilde{Q}_2(f) = -f = \widetilde{Q}'_2(f)$.*

Proof. If $f \in S^-_{k+1/2}(4p)$ then by Proposition 6.16 the conditions hold. Now assume that $f \in S_{k+1/2}(4p)$ is in the intersection of -1 eigenspaces of $\widetilde{Q}_p$, $\widetilde{Q}'_p$, $\widetilde{Q}_2$ and $\widetilde{Q}'_2$. For every $g \in S_{k+1/2}(\Gamma_0(4))$ we have $\widetilde{Q}_p(g) = pg$. Since $\widetilde{Q}_p$ is self-adjoint,

$$-\langle f, g\rangle = \langle \widetilde{Q}_p f, g\rangle = \langle f, \widetilde{Q}_p g\rangle = p\langle f, g\rangle$$

implying $\langle f, g\rangle = 0$. Thus f is orthogonal to $R \oplus S^-_{k+1/2}(4)$. For every $g \in \widetilde{W}_{p^2} S_{k+1/2}(4)$ we have $\widetilde{Q}'_p(g) = pg$ and the same argument shows that $\langle f, g\rangle = 0$ implying f is orthogonal to $\widetilde{W}_{p^2}(R \oplus S^-_{k+1/2}(4))$. Since Kohnen's plus space is the 2-eigenspace of $\widetilde{Q}'_2$, for $g \in S^{+,\mathrm{new}}_{k+1/2}(4p)$ we have $\widetilde{Q}'_2(g) = 2g$, consequently for $g \in \widetilde{W}_4 S^{+,\mathrm{new}}_{k+1/2}(4p)$ we have $\widetilde{Q}_2(g) = 2g$. Hence $\langle f, g\rangle = 0$ for such g, that is, f is orthogonal to $S^{+,\mathrm{new}}_{k+1/2}(4p) \oplus \widetilde{W}_4 S^{+,\mathrm{new}}_{k+1/2}(4p)$. It follows that $f \in S^-_{k+1/2}(4p)$. □

6.4. **Minus space for $\Gamma_0(4M)$ for M odd and square-free.** Let $M \neq 1$ be an odd and square free natural number. We write $M = p_1 p_2 \cdots p_k$. For each $i = 1, \ldots k$ define $M_i = M/p_i$. Since $S_{k+1/2}(\Gamma_0(4M_i))$ is contained in the p_i-eigenspace of $\widetilde{Q}_{p_i}$ (Corollary 4.3(5)), following the proof of Proposition 6.6 we obtain that

Proposition 6.17. $S_{k+1/2}(\Gamma_0(4M_i)) \bigcap \widetilde{W}_{p_i^2} S_{k+1/2}(\Gamma_0(4M_i)) = \{0\}$.

Corollary 6.18. *The Niwa map $\psi : S_{k+1/2}(\Gamma_0(4M)) \to S_{2k}(\Gamma_0(2M))$ maps $S_{k+1/2}(\Gamma_0(4M_i)) \oplus \widetilde{W}_{p_i^2} S_{k+1/2}(\Gamma_0(4M_i))$ isomorphically onto $S_{2k}(\Gamma_0(2M_i)) \oplus V(p_i) S_{2k}(\Gamma_0(2M_i))$.*

Let $S^{+,\mathrm{new}}_{k+1/2}(4M)$ be the new space inside the Kohnen plus subspace of $S_{k+1/2}(4M)$. Then similarly we have

Proposition 6.19. $S^{+,\mathrm{new}}_{k+1/2}(4M) \bigcap \widetilde{W}_4 S^{+,\mathrm{new}}_{k+1/2}(4M) = \{0\}$.

Corollary 6.20. *ψ maps $S^{+,\mathrm{new}}_{k+1/2}(4M) \oplus \widetilde{W}_4 S^{+,\mathrm{new}}_{k+1/2}(4M)$ isomorphically onto $S^{\mathrm{new}}_{2k}(\Gamma_0(M)) \oplus V(2) S^{\mathrm{new}}_{2k}(\Gamma_0(M))$.*

We let $B_i = S_{k+1/2}(\Gamma_0(4M_i)) \oplus \widetilde{W}_{p_i^2} S_{k+1/2}(\Gamma_0(4M_i))$, $i = 1, \ldots k$. Define

$$E = \sum_{i=1}^{k} B_i \oplus S^{+,\mathrm{new}}_{k+1/2}(4M) \oplus \widetilde{W}_4 S^{+,\mathrm{new}}_{k+1/2}(4M).$$

Proposition 6.21. *Under ψ the space E maps isomorphically onto the old space $S_{2k}^{\text{old}}(\Gamma_0(2M))$.*

Proof. This follows from Corollary 6.18 and 6.20 and from the decomposition

$$S_{2k}^{\text{old}}(\Gamma_0(2M)) = \left(\sum_{i=1}^{k} S_{2k}(\Gamma_0(2M_i)) \oplus V(p_i) S_{2k}(\Gamma_0(2M_i))\right) \oplus \left(S_{2k}^{\text{new}}(\Gamma_0(M)) \oplus V(2) S_{2k}^{\text{new}}(\Gamma_0(M))\right).$$

□

We now define the minus space to be the orthogonal complement of E,

$$S_{k+1/2}^{-}(4M) := E^{\perp}$$

Let $f \in S_{k+1/2}^{-}(4M)$ be a Hecke eigenform for all the operators T_{q^2} where q is an odd prime satisfying $(q, M) = 1$. Let $\psi(f) = F$. The proof of the following results is identical to the proofs in the previous subsections.

Proposition 6.22. *F is up to a scalar a primitive Hecke eigenform in $S_{2k}^{new}(2M)$.*

Theorem 8. *The space $S_{k+1/2}^{-}(4M)$ has a basis of eigenforms for all the operators T_{q^2} where q is an odd prime satisfying $(q, M) = 1$. Under ψ, the space $S_{k+1/2}^{-}(4M)$ maps isomorphically onto the space $S_{2k}^{\text{new}}(\Gamma_0(2M))$. If two forms in $S_{k+1/2}^{-}(4M)$ have the same eigenvalues for all the operators T_{q^2}, $(q, 2M) = 1$, then they are same up to a scalar factor.*

Let $f \in S_{k+1/2}^{-}(4M)$ be a Hecke eigenform for all the operators T_{q^2}, $(q, 2M) = 1$. Then $\psi(f) = F$ is a Hecke eigenform in $S_{2k}^{\text{new}}(\Gamma_0(2M))$ for all operators T_q, $(q, 2M) = 1$. By [1], for all primes p such that $p|M$, $U_p(F) = -p^{k-1}\lambda(p)F$ where $\lambda(p) = \pm 1$ and $U_2(F) = -2^{k-1}\lambda(2)F$ where $\lambda(2) = \pm 1$.

Proposition 6.23. *Let $f \in S_{k+1/2}^{-}(4M)$ be a Hecke eigenform for all the operators T_{q^2}, q prime, $(q, 2M) = 1$. Then for all primes p such that $p|M$*

$$U_{p^2}(f) = -p^{k-1}\lambda(p)f \quad \text{and} \quad U_4(f) = -2^{k-1}\lambda(2)f$$

where $\lambda(p) = \pm 1$ and $\lambda(2) = \pm 1$ are defined as above.

Following [10, Theorem 1.9] we have

Corollary 6.24. *Let $f = \sum_{n=0}^{\infty} a_n q^n \in S_{k+1/2}^{-}(4M)$ be a Hecke eigenform for all Hecke operators, i.e, $T_{q^2}(f) = \omega_q f$ for all primes $(q, 2M) = 1$ and $U_{p^2}(f) = \omega_p f$ for all primes $p \mid 2M$. Let $F = \sum_{n=0}^{\infty} A_n q^n \in S_{2k}^{\text{new}}(\Gamma_0(2M))$ be the unique normalized primitive form determined by f, i.e. $A_p = \omega_p$ for all primes p. Then for a fundamental discriminant D such that $(-1)^k D > 0$,*

$$\mathrm{L}\left(s - k + 1, \left(\frac{D}{\cdot}\right)\right) \sum_{n=1}^{\infty} a_{|D|n^2}\ n^{-s} = a(|D|) \sum_{n=1}^{\infty} A_n n^{-s}.$$

We finally give the characterization of our minus space. The proofs of the following proposition and theorem are as before.

Proposition 6.25. *Let $f \in S^-_{k+1/2}(4M)$. Then for every prime p dividing M we have $\widetilde{Q}_p(f) = -f = \widetilde{Q}'_p(f)$ and $\widetilde{Q}_2(f) = -f = \widetilde{Q}'_2(f)$.*

Theorem 9. *Let $f \in S_{k+1/2}(4M)$. Then $f \in S^-_{k+1/2}(4M)$ if and only if $\widetilde{Q}_p(f) = -f = \widetilde{Q}'_p(f)$ for every prime p dividing M and $\widetilde{Q}_2(f) = -f = \widetilde{Q}'_2(f)$.*

6.5. **Some examples.** We complete this section by giving two examples.

Example 1. The space $S_{3/2}(\Gamma_0(28))$ is one dimensional and is spanned by

$$f = q - q^2 - q^4 + q^7 + q^8 - q^9 + q^{14} - 2q^{15} + q^{16} + 3q^{18} - 2q^{21} + \dots$$

By Shimura decomposition (for notation see [9])

$$S_{3/2}(\Gamma_0(28)) = \bigoplus_{\substack{F \in S_2^{\text{new}}(\Gamma_0(M)) \\ \text{prim., } M|14}} S_{3/2}(28, F) = S_{3/2}(28, F_{14})$$

since there are no primitive forms of weight 2 at level 1, 2, 7 and $F_{14} \in S_2^{\text{new}}(\Gamma_0(14))$ is the only primitive form at level 14. In particular, we have $S^+_{3/2}(28) = \{0\}$ and $S^-_{3/2}(28) = S_{3/2}(\Gamma_0(28)) = \langle f \rangle$.

Example 2. The space $S_{7/2}(\Gamma_0(12))$ is three dimensional and is spanned by

$$g_1 = q - 4q^4 - 6q^6 + 8q^7 + 9q^9 + 4q^{10} + 12q^{12} - 20q^{13} - 24q^{15} + \dots$$
$$g_2 = q^3 - 2q^4 + 2q^7 - 2q^{12} - 6q^{15} + 12q^{16} - 10q^{19} + 12q^{24} + \dots$$
$$g_3 = q^2 - q^3 - 4q^5 + 3q^6 + 4q^8 + 2q^{11} - 4q^{12} - 8q^{14} + 16q^{17} + \dots$$

We have two primitive forms of weight 6 and level dividing 6, namely F_3 of level 3 and F_6 of level 6. Using algorithm in [9] we have

$$S_{7/2}(\Gamma_0(12)) = S_{7/2}(12, F_3) \oplus S_{7/2}(12, F_6) = \langle g_1,\ g_2 \rangle \oplus \langle g_3 \rangle.$$

Note that g_2 is in the plus space, so we have $S^+_{7/2}(12) = \langle g_2 \rangle$. The minus space $S^-_{7/2}(12) = S_{7/2}(12, F_6) = \langle g_3 \rangle$.

Remark 5. *(i) In general, $S^-_{k+1/2}(4M) = \bigoplus_F S_{k+1/2}(4M, F)$ where F runs through all primitive forms of weight $2k$ and level $2M$.*
(ii) The Kohnen plus space is given by well-known Fourier coefficient condition. But we do not expect any such Fourier coefficient condition for forms in our minus space. This is also evident from the above examples.

Appendix A. Some observations on cocycle multiplication

Let p denotes any prime. In this appendix we note down some useful observations on multiplication in $\widetilde{\mathrm{SL}}_2(Q_p)$ by cocycle σ_p.

Recall the Hilbert symbol $(\cdot,\cdot)_p$ defined on $\mathbb{Q}_p^\times \times \mathbb{Q}_p^\times$. For an odd prime p it can be given by the formula: For a, b coprime to p,

$$\left(p^s a, p^t b\right)_p = \left(\frac{-1}{p}\right)^{st} \left(\frac{a}{p}\right)^t \left(\frac{b}{p}\right)^s.$$

Thus $(p,p)_p = \left(\frac{-1}{p}\right)$ and $(-p,u)_p = (p,u)_p = \left(\frac{u}{p}\right)$ where u is a unit in $\mathbb{Z}_p$. For an even prime, if a, b are odd

$$\left(2^s a, 2^t b\right)_2 = (-1)^{\frac{(a-1)(b-1)}{4}} \left(\frac{2}{|a|}\right)^t \left(\frac{2}{|b|}\right)^s.$$

Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Q}_p)$. For $(A,\epsilon_1) \in \widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$, $(A,\ \epsilon_1)^{-1} = (A^{-1},\ \epsilon_1\sigma_p(A,A^{-1}))$ where

(i) If $c=0$ then $\sigma_p(A,A^{-1}) = (a,a)_p = (d,d)_p$.
(ii) If $c\neq 0$ and $\mathrm{ord}_p(c)$ is even then $\sigma_p(A,A^{-1}) = 1$.
(iii) If $c\neq 0$ and $\mathrm{ord}_p(c)$ is odd then

$$\sigma_p(A,A^{-1}) = \begin{cases} (c,d)_p\,(-c,a)_p & \text{if } d\neq 0,\ a\neq 0 \\ (c,d)_p & \text{if } d\neq 0,\ a=0 \\ (-c,a)_p & \text{if } d=0,\ a\neq 0 \\ 1 & \text{if } d=0,\ a=0. \end{cases}$$

In particular if $A \in \{x(p^n),\ y(p^n),\ w(p^n)\}_{n\in\mathbb{Z}}$ then $\sigma_p(A,A^{-1}) = 1$. For $A = h(p^n)$ with $n\in\mathbb{Z}$, if $p=2$ then $\sigma_p(A,A^{-1}) = 1$, however if p is an odd prime then

$$\sigma_p(A,A^{-1}) = \begin{cases} 1 & \text{if } n \text{ even}, \\ \left(\frac{-1}{p}\right) & \text{else.} \end{cases}$$

Let $(A,\epsilon_1),\ (B,\ \epsilon_2) \in \widetilde{\mathrm{SL}}_2(\mathbb{Q}_p)$. The following lemmas can be easily obtained using cocycle formula.

Lemma A.1. *We have* $[(B,\ \epsilon_2)^{-1},\ (A,\ \epsilon_1)^{-1}] = (B^{-1}A^{-1}BA, \xi)$ *where* $\xi = \sigma_p(A,A^{-1})\sigma_p(B,B^{-1})\sigma_p(B,A)\sigma_p(A^{-1},BA)\sigma_p(B^{-1},A^{-1}BA)$.

Lemma A.2. *The* σ_p*-factor (*ξ *factor above) of* $[(B,\ \epsilon_2)^{-1},\ (A,\ \epsilon_1)^{-1}]$ *equals*

$$\Big(\tau(B),\ \tau(B^{-1})\Big)_p \Big(\tau(A),\ \tau(A^{-1})\Big)_p$$
$$\Big(\tau(BA)\tau(B),\ \tau(BA)\tau(A)\Big)_p \Big(\tau(A^{-1}BA)\tau(A^{-1}),\ \tau(A^{-1}BA)\tau(BA)\Big)_p$$
$$\Big(\tau(B^{-1}A^{-1}BA)\tau(B^{-1}),\ \tau(B^{-1}A^{-1}BA)\tau(A^{-1}BA)\Big)_p\, s_p(B^{-1}A^{-1}BA).$$

In the proofs for checking support of our local Hecke algebra (section 3) we required the following lemma.

Lemma A.3. *Let* $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Q}_p)$. *Then*

(a) If $B = x(s)$ *where* $s \neq 0$, *then* σ_p*-factor is*

$$\begin{cases} \left(-sc^2,\ 1-cds\right)_p & \text{if } sc^2(1-cds) \neq 0 \text{ and } \mathrm{ord}_p(s) \text{ is odd,} \\ 1 & \text{else.} \end{cases}$$

(b) If $B = h(u)$ *where* $u \neq \pm 1$, *then* σ_p*-factor is*

$$\begin{cases} \left(ac(1-u^2),\ 1+(1-u^2)bc\right)_p & \text{if } ac(1-u^2)(1+(1-u^2)bc) \neq 0 \\ & \text{and } \mathrm{ord}_p(ac(1-u^2)) \text{ is odd,} \\ 1 & \text{else.} \end{cases}$$

(c) If $B = y(t)$ *where* $t \neq 0$, *then* σ_p*-factor is*

$$\begin{cases} \left((a^2-1)t+abt^2, 1+abt+b^2t^2\right)_p & \text{if } ((a^2-1)t+abt^2)(1+abt+b^2t^2) \neq 0 \\ & \text{and } \mathrm{ord}_p((a^2-1)t+abt^2) \text{ is odd,} \\ 1 & \text{else.} \end{cases}$$

In each of the above cases the σ_p*-factor is simply* $s_p(B^{-1}A^{-1}BA)$.

Proof. For (a) let $B = x(s)$ where $s \neq 0$. Then we have

$$BA = \begin{pmatrix} a+sc & b+sd \\ c & d \end{pmatrix}, \quad A^{-1}BA = \begin{pmatrix} 1+cds & sd^2 \\ -sc^2 & 1-cds \end{pmatrix},$$

$$B^{-1}A^{-1}BA = \begin{pmatrix} 1+cds+s^2c^2 & sd^2-s+cds^2 \\ -sc^2 & 1-cds \end{pmatrix}.$$

It is easy to see that $\left(\tau(B),\ \tau(B^{-1})\right)_p = 1$ and that $\left(\tau(A),\ \tau(A^{-1})\right)_p =$

$$= \left(\tau(A^{-1}BA)\tau(A^{-1}),\ \tau(A^{-1}BA)\tau(BA)\right)_p = \begin{cases} 1 & \text{if } c \neq 0 \\ (d,a)_p & \text{else.} \end{cases}$$

Further one can check that $\left(\tau(BA)\tau(B), \tau(BA)\tau(A)\right)_p = 1$ and also

$$\left(\tau(B^{-1}A^{-1}BA)\tau(B^{-1}),\ \tau(B^{-1}A^{-1}BA)\tau(A^{-1}BA)\right)_p = 1.$$

Finally we have $s_p(B^{-1}A^{-1}BA)$

$$= \begin{cases} \left(-sc^2,\ 1-cds\right)_p & \text{if } sc^2(1-cds) \neq 0 \text{ and } ord_p(s) \text{ is odd,} \\ 1 & \text{else.} \end{cases}$$

By using Lemma A.2, multiplying all the above terms we get the required σ_p-factor.

For (b) we proceed similarly. Let $B = h(u)$ where $u \neq \pm 1$. Then

$$BA = \begin{pmatrix} ua & ub \\ u^{-1}c & u^{-1}d \end{pmatrix}, \quad A^{-1}BA = \begin{pmatrix} uad - u^{-1}bc & bd(u - u^{-1}) \\ ac(u^{-1} - u) & u^{-1}ad - ubc \end{pmatrix},$$

$$B^{-1}A^{-1}BA = \begin{pmatrix} 1 + (1 - u^{-2})bc & bd(1 - u^{-2}) \\ ac(1 - u^2) & 1 + (1 - u^2)bc \end{pmatrix}.$$

We have $\Big(\tau(B),\ \tau(B^{-1})\Big)_p = \big(u, u^{-1}\big)_p$. Also $\Big(\tau(A),\ \tau(A^{-1})\Big)_p = 1$ if $c \neq 0$ and $(d, a)_p$ else. We check that

$$\Big(\tau(BA)\tau(B),\ \tau(BA)\tau(A)\Big)_p = \begin{cases} \big(c, u^{-1}\big)_p & \text{if } c \neq 0 \\ \big(d, u^{-1}\big)_p & \text{else,} \end{cases}$$

$$\Big(\tau(A^{-1}BA)\tau(A^{-1}),\ \tau(A^{-1}BA)\tau(BA)\Big)_p$$

$$= \begin{cases} \big(-a(u^{-1} - u), u^{-1}\big)_p & \text{if } ac \neq 0 \\ (bu, -b)_p & \text{if } a = 0 \text{ and } c \neq 0 \\ \big(du^{-1}, a\big)_p & \text{if } a \neq 0 \text{ and } c = 0, \end{cases}$$

$$\Big(\tau(B^{-1}A^{-1}BA)\tau(B^{-1}),\ \tau(B^{-1}A^{-1}BA)\tau(A^{-1}BA)\Big)_p$$

$$= \begin{cases} \big(ac(u^{-1} - u), u^{-1}\big)_p & \text{if } ac \neq 0 \\ (bc, u)_p = (1, u)_p & \text{if } a = 0 \text{ and } c \neq 0 \\ (-ad, u)_p & \text{if } a \neq 0 \text{ and } c = 0, \end{cases}$$

and $s_p(B^{-1}A^{-1}BA) =$

$$\begin{cases} \big(ac(1 - u^2),\ 1 + (1 - u^2)bc\big)_p & \text{if } ac(1-u^2)(1+(1-u^2)bc) \neq 0 \\ & \text{and } \mathrm{ord}_p(ac(1 - u^2)) \text{ is odd,} \\ 1 & \text{else.} \end{cases}$$

Again by multiplying all the above terms we get the required σ_p-factor.

For (c), let $B = y(t)$ where $t \neq 0$. Then

$$BA = \begin{pmatrix} a & b \\ at + c & bt + d \end{pmatrix}, \quad A^{-1}BA = \begin{pmatrix} 1 - abt & -b^2t \\ a^2t & 1 + abt \end{pmatrix},$$

$$B^{-1}A^{-1}BA = \begin{pmatrix} 1 - abt & -b^2t \\ (a^2 - 1)t + abt^2 & 1 + abt + b^2t^2 \end{pmatrix}.$$

As before, $\Big(\tau(B),\ \tau(B^{-1})\Big)_p = (t, -t)_p = 1$, and $\Big(\tau(A),\ \tau(A^{-1})\Big)_p = 1$ if $c \neq 0$ and $(d, a)_p$ else. One can compute (using $ad - bc = 1$ in the Hilbert symbol calculations) that

$$\Big(\tau(BA)\tau(B),\ \tau(BA)\tau(A)\Big)_p = \begin{cases} (t(at + c), -ct)_p & \text{if } a \neq -c/t \text{ and } c \neq 0 \\ (-c, a)_p & \text{if } a = -c/t \text{ and } c \neq 0 \\ (a, -dt)_p & \text{if } c = 0, \end{cases}$$

$$\Big(\tau(A^{-1}BA)\tau(A^{-1}),\ \tau(A^{-1}BA)\tau(BA)\Big)_p =$$

$$= \begin{cases} (t(at+c), -ct)_p & \text{if } a \neq -c/t \text{ and } c \neq 0 \text{ and } a \neq 0 \\ 1 & \text{if } a \neq -c/t \text{ and } c \neq 0 \text{ and } a = 0 \\ (-c, a)_p & \text{if } a = -c/t \text{ and } c \neq 0 \\ (a, at)_p & \text{if } c = 0. \end{cases}$$

All the above factors clearly multiplies to 1. Also it turns out that

$$\Big(\tau(B^{-1}A^{-1}BA)\tau(B^{-1}),\ \tau(B^{-1}A^{-1}BA)\tau(A^{-1}BA)\Big)_p = 1,$$

so we get the required σ_p-factor. □

We also note the triangular decomposition of $\overline{K_0^p(p^n)}$.

Lemma A.4. *We have a triangular decomposition*

$$\overline{K_0^p(p^n)} = N^{\overline{K_0^p(p^n)}} T^{\overline{K_0^p(p^n)}} \bar{N}^{\overline{K_0^p(p^n)}}.$$

More precisely for $(A, \epsilon) = (\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \epsilon) \in \overline{K_0^p(p^n)}$,

$$(A, \epsilon) = (x(s), 1)(h(u), 1)(y(t), 1)(I, \epsilon\delta)$$

where

$$u = d^{-1},\ s = d^{-1}b,\ t = d^{-1}c,$$

and

$$\delta = \begin{cases} 1 & c = 0 \\ (d, -1)_p & c \neq 0,\ \mathrm{ord}_p(c) \textit{ is odd} \\ (-c, d)_p & c \neq 0,\ \mathrm{ord}_p(c) \textit{ is even.} \end{cases}$$

Proof. Clearly

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & bd^{-1} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} d^{-1} & 0 \\ 0 & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ cd^{-1} & 1 \end{pmatrix}.$$

Let $u = d^{-1},\ s = bd^{-1},\ t = cd^{-1}$. Since

$$x(s)h(u)y(t) = \begin{pmatrix} u & su^{-1} \\ 0 & u^{-1} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix} = \begin{pmatrix} u + su^{-1}t & su^{-1} \\ tu^{-1} & u^{-1} \end{pmatrix},$$

we get that

$$(x(s), 1)(h(u), 1)(y(t), 1) = (x(s)h(u)y(t), \delta) = (A, \delta)$$

where

$$\delta = \sigma(x(s), h(u))\sigma(x(s)h(u), y(t)) = \begin{cases} 1 & t = 0 \\ (u, -1)_p & t \neq 0,\ \mathrm{ord}_p(t) \text{ is odd} \\ (t, u)_p & t \neq 0,\ \mathrm{ord}_p(t) \text{ is even.} \end{cases}$$

Substituting u, s, t in terms of b, c, d we get δ as in the statement. □

References

[1] A. O. L. Atkin and J. Lehner, *Hecke operators on* $\Gamma_0(m)$, Math. Ann. **185** (1970), 134–160.
[2] E. M. Baruch and S. Purkait, *Hecke algebras, new vectors and newforms on* $\Gamma_0(m)$, on arXiv.
[3] S. Gelbart, *Weil's representation and the spectrum of the metaplectic group*, Lecture Notes in Math. **530**, Springer, Berlin, 1976.
[4] W. Kohnen, *Modular forms of half-integral weight on* $\Gamma_0(4)$, Math. Ann. **248** (1980), 249–266.
[5] W. Kohnen, *Newforms of half-integral weight*, J. Reine Angew. Math. **333** (1982), 32–72.
[6] H. Y. Loke and G. Savin, *Representations of the two-fold central extension of* $\mathrm{SL}_2(\mathbb{Q}_2)$, Pacific J. Math. **247** (2010), 435–454.
[7] S. Niwa, *On Shimura's trace formula*, Nagoya Math. J. **66** (1977), 183–202.
[8] S. Purkait, *On Shimura's decomposition*, Int. J. Number Theory **9** (2013), 1431-1445.
[9] S. Purkait, *Hecke operators in half-integral weight*, J. Théor. Nombres Bordeaux **26** (2014), 233–251.
[10] G. Shimura, *On modular forms of half integral weight*, Ann. of Math. **97** (1973), 440–481.
[11] M. Ueda, *On twisting operators and newforms of half-integral weight*, Nagoya Math J. **131** (1993), 135–205.
[12] J.-L. Waldspurger, *Sur les coefficients de Fourier des formes modulaires de poids demi-entier*, J. Math. Pures Appl. (9) **60** (1981), 375–484.

Department of Mathematics, Technion, Haifa , 32000, Israel
E-mail address: embaruch@math.technion.ac.il

Faculty of Mathematics, Kyushu University, Japan
E-mail address: somapurkait@gmail.com